arXiv:1004.0359v1 [cond-mat.quant-gas] 2 Apr 2010

Many Body Approach for Quartet Condensation in Strong Coupling.

Takaaki Sogo, Gerd Röpke
Institut für Physik, Universität Rostock, D-18051 Rostock, Germany

Peter Schuck
Institut de Physique Nucléaire, CNRS, UMR 8608, Orsay F-91406, France
Université Paris-Sud, Orsay F-91505, France
Laboratoire de Physique et Modélisation des Milieux Condensés,
CNRS and Université Joseph Fourier, 25 Avenue des Martyrs,
Boîte Postale 166, F-38042 Grenoble Cedex 9, France and
Groupe de Physique Theorique Institut de Physique Nucleaire, 91406 Orsay Cedex, France

The theory for condensation of higher fermionic clusters is developed. Fully selfconsistent nonlinear equations for the quartet order parameter in strongly coupled fermionic systems are established and solved. The breakdown of the quasiparticle picture is pointed out. Derivation of numerically tractable approximation is described. The momentum projected factorisation ansatz of Ref. [21] for the order parameter is employed again. As a definite example the condensation of α particles in nuclear matter is worked out.

I. INTRODUCTION

Quartet condensation is relevant in several domains of physics. It is so far mostly considered in nuclear physics with the strongly bound alpha particle cluster playing a dominant role in certain states of lighter nuclei [1] and, eventually, also in the surface of heavy nuclei as may be indicated by observed alpha decay processes. Alpha particle condensation may eventually occur in compact stars [2]. However, trapping of multi component fermionic atoms makes fast progress. The case of trions is already quite advanced, experimentally [3, 4] and also with theory [5–13]. One may trap fermions with four different 'colors' in the near future. This is a prerequisite for quartet formation and quartet condensation. Theoretical work on this subject already has appeared [14–18]. The condensation of bi- excitons in semi-conductors also may be of relevance [19, 20].

In the recent past we already have published several papers on alpha particle condensation in infinite nuclear matter. For instance, we studied the onset of alpha particle condensation and evaluated the corresponding critical temperature [21, 22] with a procedure analogous to the pairing case via a four body generalization of the famous Thouless criterion [23]. However, quartet condensation not only is formally much more complicated than condensation of (Cooper) pairs, it turns out that also certain aspects of the physics are quite different. The most striking feature being that quartets only exist in the so-called BEC (Bose-Einstein condensate) limit where they do not overlap very much in space. Contrary to the pairing case quartets cannot strongly intermingle in real space and, therefore, a coherence length much longer than the inter alpha particle distance cannot exist. We will give the reason for this different behavior in the main text.

In this paper we will treat quartet condensation at zero temperature and establish and solve a full nonlinear equation for the quartet order parameter which will be the analog to the gap equation for pairing. A direct solution of such a highly non linear four body problem seems hopeless, even in homogeneous matter. However, we recently showed that a very simplifying approximation works very well, at least in the domain of negative chemical potentials, i.e. in the strong coupling regime. This approximation consists in making a mean field ansatz, i.e. a Slater determinant, for the quartet but projected onto zero total momentum as it is relevant for condensation [21]. Such a mean field treatment may work each time the quartet is in its lowest energy configuration. However, one may think of generalisations for excited configurations as well. Since the four fermions of the quartet are all different in the nuclear case (proton/neutron, spin up/down), all are in the same 0S mean field wave function and the problem boils down in the end to solve the equation in iterating on this single one particle mean field wave function. The same happens, of course, if the four fermions are components of an $F = 3/2$ spin as in [16]. The problem still is complicated but can be solved with effective interactions of separable form. It was shown that that approximation gives comparable results as with the four body Faddeev-Yakubovsky equation with a more realistic interaction in the nuclear case. So, in the present work we calculate the order parameter equation with the projected mean field ansatz and a separable potential.

The paper is organized as follows: in the next section, we discuss the BCS gap equation. In Sec. III, we show the full expression of the single particle mass operator for quartet condensation. Since this is too difficult to calculate numerically, we suggest an approximate mass operator in Sec. IV. Before we show the numerical result employed by the approximated mass operator, we discuss in Sec. V the significant difference between pairing condensation and quartet condensation through the different level densities involved in the condensation processes. In Sec. VI, we present the results. Finally we conclude in the last section. In the Appendix, we describe the detailed derivation of the equations and discuss various methods

to formulate the order parameter equation for quartet condensation.

II. RECAPITULATION OF THE PAIRING CASE

In order to prepare the terrain for our procedure in the quartetting case, in this section we want to rederive standard BCS theory in a way somewhat differing from the usual.

The one body Green's function (GF) for BCS is represented by [26]

$$G_{1;1'}(\omega) = \frac{\delta_{11'}}{\omega - \varepsilon_1 - M^{\rm BCS}_{1;1}(\omega)}, \tag{1}$$

where $M^{\rm BCS}_{1;1}(\omega)$ is the BCS mass operator

$$M^{\rm BCS}_{1;1'}(\omega) = \sum_2 \frac{\Delta_{12}\Delta^*_{1'2}}{\omega + \varepsilon_2}, \tag{2}$$

where

$$\Delta_{12} = -\frac{1}{2}\sum_{34} \bar{v}_{12,34}\langle c_4 c_3\rangle \tag{3}$$

with $\langle cc\rangle$ being the thermal average of the pair operator.

The indices $1, 2, 3, ...$ correspond to momentum and spin. In nuclear matter, we have to add isospin. $\bar{v}_{12,34}$ is the antisymmetrized matrix element of the two-body interaction ($\bar{v}_{12,34} = -\bar{v}_{12,43} = -\bar{v}_{21,34}$). The single particle energies ε_i are in principle given by the kinetic energies plus the mean field shifts. The direct term is in homogeneous matter a constant which can be incorporated into the chemical potential and the Fock-term gives rise to an effective mass. Since we will mostly deal with very low density nuclear matter, we will not consider a mass renormalization here. Therefore, in Eq. (1) we have $\varepsilon_1 = k_1^2/(2m) - \mu_1$ with the chemical potential μ_1 which, in principle, contains the direct part of the mean field. We have attached an index on the chemical potential, since in principal it can depend on the various fermionic components involved. However, in this work we always will consider fully symmetric situations and henceforth we will suppose that the chemical potentials of all fermionic species are equal and, therefore, drop the index.

Fig. 1 is the graphical representation of the BCS mass operator. As shown in Eq. (2) and Figure 1, the BCS mass operator consists of the 2 particle-1 hole (2p1h) GF between two-body vertices factorized into two order parameters and the free 1h GF.

On the other hand, the in-medium Schrödinger equation for the order parameter is of the following form

$$\langle c_2 c_1\rangle = -\frac{1-\rho_1-\rho_2}{\varepsilon_1+\varepsilon_2}\sum_{1'2'}\frac{1}{2}\bar{v}_{12,1'2'}\langle c_{2'}c_{1'}\rangle, \tag{4}$$



FIG. 1: Graphic representation of the BCS mass operator in Eq. (2)



FIG. 2: Some diagram for the mass operator.

where ρ_1 is the occupation number derived from

$$\rho_1 = -\int \frac{d\omega}{2\pi} 2{\rm Im}G_{1;1}(\omega + i\eta) f(\omega) \tag{5}$$

with the Fermi distribution function $f(\omega) = [e^{\omega/T}+1]^{-1}$ and a positive infinitesimal of η, as indicated.

In the standard BCS theory, pairs in time reversed states are considered, i.e. in Eqs. (3) and (4) taking $2 = \bar{1}$, $\rho_1 = \rho_{\bar{1}}$, and $\varepsilon_1 = \varepsilon_{\bar{1}}$, where $\bar{1}$ is the time reversal conjugate of quantum numbers 1. For Eq. (1), we obtain the imaginary part of the one body Green function as

$$\begin{aligned} -{\rm Im}G_{1;1}(\omega+i\eta) &= \frac{1}{2}\Big(1+\frac{\varepsilon_1}{E_1}\Big)\pi\delta(\omega - E_1) \\ &+ \frac{1}{2}\Big(1-\frac{\varepsilon_1}{E_1}\Big)\pi\delta(\omega + E_1) \end{aligned} \tag{6}$$

with $\Delta_{1\bar{1}} = \Delta_1$ and $E_1 = \sqrt{\varepsilon_1^2 + \Delta_1^2}$. This is equivalent to solving the usual gap equation at finite temperature as can easily be deduced from the spectral function obtained from (1) [24, 25]:

$$\Delta_1 = -\sum_{1'}\frac{1}{2}\bar{v}_{1\bar{1},1'\bar{1}'}\frac{\Delta_{1'}}{2E_{1'}}\tanh(\frac{E_{1'}}{2T}). \tag{7}$$

Note that Eq. (4) resembles a particle-particle RPA equation [26] with, however, renormalized occupation numbers. One could, therefore, also consider Eq. (4) as a single pole approximation to the so-called renormalized RPA, well known in the literature, see, e.g., Ref. [27].

III. SINGLE PARTICLE MASS OPERATOR AND QUARTET CONDENSATION

Formally it is straightforward to generalize the pairing case to the quartet case. A typical diagram of the

mass operator needed for quartet condensation is shown in Fig. 2. As seen, to make the quartet order parameter appear in the single particle mass operator, we need to express it by the 4p3h GF. In order to get to it, we must successively open phase space from one particle to 2p1h, to 3p2h, to 4p3h. This needs three interaction vertices on each side as shown in Fig. 2. Details of the derivation will be given in the Appendix B. We obtain for the 1p mass operator with quartet condensation

$$\begin{aligned} & M^{\rm quartet}_{1;1'}(\omega) \\ = \; & \Gamma^{(4)}_{1234;5678}(\omega)\langle c_8 c_7 c_6 c_5\rangle \frac{\bar f_2\bar f_3\bar f_4 + f_2 f_3 f_4}{\omega+\varepsilon_{234}} P_{234;2'3'4'} \\ & \times\langle c^\dagger_{5'} c^\dagger_{6'} c^\dagger_{7'} c^\dagger_{8'}\rangle \Gamma^{(4)*}_{1'2'3'4';5'6'7'8'}(\omega), \end{aligned} \tag{8}$$

where summation convention over repeated indices is to be understood, and

$$P_{123;1'2'3'} = \begin{vmatrix} \delta_{11'} & \delta_{12'} & \delta_{13'} \\ \delta_{21'} & \delta_{22'} & \delta_{23'} \\ \delta_{31'} & \delta_{32'} & \delta_{33'} \end{vmatrix}. \tag{9}$$

The effective four body vertex $\Gamma^{(4)}$, evaluated perturbatively to third order in the interaction is

$$\begin{aligned} & \Gamma^{(4)}_{1234;5678}(\omega) \\ = \; & \Gamma^{(3)}_{12'3';4'5'6'}(\omega)\frac{1}{\omega-\varepsilon_{4'5'6'}+\varepsilon_{2'3'}} \\ \times \; & \Big[\frac{1}{2}\bar v_{4'4,58}\delta_{2'2}\delta_{3'3}\delta_{5'6}\delta_{6'7} + \frac{1}{2}\bar v_{5'4,68}\delta_{2'2}\delta_{3'3}\delta_{4'5}\delta_{6'7} \\ & + \frac{1}{2}\bar v_{6'4,78}\delta_{2'2}\delta_{3'3}\delta_{4'5}\delta_{5'6} - \frac{1}{2}\bar v_{2'8,24}\delta_{3'3}\delta_{4'5}\delta_{5'6}\delta_{6'7} \\ & - \frac{1}{2}\bar v_{3'8,34}\delta_{2'2}\delta_{4'5}\delta_{5'6}\delta_{6'7}\Big], \end{aligned} \tag{10}$$

$$\begin{aligned} & \Gamma^{(3)}_{123;456}(\omega) \\ = \; & \frac{1}{2}\bar v_{12',3'4'}\frac{1}{\omega-\varepsilon_{3'4'}+\varepsilon_{2'}}\Big[\frac{1}{2}\bar v_{3'2,45}\delta_{2'3}\delta_{4'6} \\ + \; & \frac{1}{2}\bar v_{4'3,45}\delta_{2'2}\delta_{3'6} - \frac{1}{2}\bar v_{2'6,23}\delta_{3'4}\delta_{4'5}\Big]. \end{aligned} \tag{11}$$

where $\varepsilon_{123\ldots}=\varepsilon_1+\varepsilon_2+\varepsilon_3+\cdots$, and $\bar f_i = 1-f_i$ with the Fermi distribution function $f_i = f(\varepsilon_i)$.

One may wonder why in (10) and (11) no Fermi function factors appear with the propagators. This, however, is a general feature of vertices coupling lower configuration spaces to higher ones. A fameous example in the literature is given by coupling 1p1h and 2p2h spaces as this appears in the damping of zero sound modes, see, e.g. [29]. One may notice that the absence of Pauli blocking factors opens up phase space and, therefore, enhances the coupling.

One can ask the question whether in Eq. (8) the uncorrelated mean field 3h GF should be used. One may think to include the hole GF's into the selfconsistent cycle or even include higher correlations. One should notice, however, that in BCS this is not done and in Eq. (2) the mean field 1h GF is used. This has a good reason, since BCS theory is based on a variational wave function which fully respects the Pauli principle. Should a self consistent hole GF be used in (2), this property would be lost. We, therefore, also stick to the mean field hole GF's in the quartet case.

We also need the in-medium four-body Schrödinger equation for the order parameter, in analogy with the pairing case shown in Eq. (4) in the previous section. It is given by

$$\begin{aligned} & \varepsilon_{1234}\langle c_4 c_3 c_2 c_1\rangle \\ & + \sum_{1'2'3'4'} V_{1234;1'2'3'4'}\langle c_{4'} c_{3'} c_{2'} c_{1'}\rangle = 0, \end{aligned} \tag{12}$$

where

$$\begin{aligned} & V_{1234;1'2'3'4'} \\ = \; & (1-\rho_1-\rho_2)\frac{1}{2}\bar v_{12;1'2'}\delta_{33'}\delta_{44'} \\ + \; & (1-\rho_1-\rho_3)\frac{1}{2}\bar v_{13;1'3'}\delta_{22'}\delta_{44'} \\ + \; & \text{permutations}. \end{aligned} \tag{13}$$

Details are given in [22] and Appendix C.

We consider symmetric (nuclear) matter. In this case, we can give a fully symmetric order parameter of exchange between two particles with respect to momenta:

$$\langle c_4 c_3 c_2 c_1\rangle \to \phi_{\vec k_1,\vec k_2,\vec k_3,\vec k_4}\chi_0, \tag{14}$$

where the spin-isospin singlet wave function is represented by χ_0, and we consider here a spin-isospin independent two-body interaction:

$$\begin{aligned} \bar v_{12,34} \; \to \; & v_{\vec k_1\vec k_2,\vec k_3\vec k_4}(\delta_{s_1s_3}\delta_{t_1t_3}\delta_{s_2s_4}\delta_{t_2t_4} \\ & -\delta_{s_1s_4}\delta_{t_1t_4}\delta_{s_2s_3}\delta_{t_2t_3}) \end{aligned} \tag{15}$$

with s_i (t_i) of spin (isospin) index. $v_{\vec k_1\vec k_2,\vec k_3\vec k_4}$ is symmetric with respect to exchange of the momenta: $v_{\vec k_1\vec k_2,\vec k_3\vec k_4} = v_{\vec k_2\vec k_1,\vec k_3\vec k_4} = v_{\vec k_1\vec k_2,\vec k_4\vec k_3}$.

Then, Eq. (12) is explicitly written as

$$\begin{aligned} & \sum_{i=1}^4 \varepsilon_i \phi_{\vec k_1\vec k_2\vec k_3\vec k_4} + \int\prod_{i=1}^4 \frac{d^3k_i'}{(2\pi)^3} \\ \times \; & \Big[(1-\rho(\vec k_1)-\rho(\vec k_2))v_{\vec k_1\vec k_2,\vec k_1'\vec k_2'} \\ & \quad\times(2\pi)^3\delta(\vec k_3-\vec k_3')(2\pi)^3\delta(\vec k_4-\vec k_4') \\ & +(1-\rho(\vec k_1)-\rho(\vec k_3))v_{\vec k_1\vec k_3,\vec k_1'\vec k_3'} \\ & \quad\times(2\pi)^3\delta(\vec k_2-\vec k_2')(2\pi)^3\delta(\vec k_4-\vec k_4') \\ & +\text{permutations}\Big]\phi_{\vec k_1'\vec k_2'\vec k_3'\vec k_4'} = 0. \end{aligned} \tag{16}$$

A sketch of the quartet mass operator is shown in Fig. 3.



FIG. 3: Graphic representation of the mass operator for alpha condensation in Eq. (8).



FIG. 4: The graphical representation of the approximate α-BEC mass operator $M^{\rm quartet}$ of Eq. (17).

IV. APPROXIMATE QUARTET MASS OPERATOR

From Eqs. (10) and (11) it becomes evident that the full evaluation of the 1p $\leftrightarrow$ 4p3h vertices are too complicated to be evaluated exactly. However, with quite reasonable approximations, one arrives at a numerically manageable expression. Since this discussion involves lengthy, quite technical details, we relegate it to Appendix D and only give the final result here:

$$M^{\rm quartet}_{1;1}(\omega) = \sum_{234} \frac{\tilde{\Delta}_{1234}(\bar{f}_2\bar{f}_3\bar{f}_4 + f_2f_3f_4)\tilde{\Delta}^*_{1234}}{\omega + \varepsilon_{234}} \quad (17)$$

where the quartet 'gap' matrix $\tilde{\Delta}_{1234}$ is given by

$$\tilde{\Delta}_{1234} = \lambda' \frac{1}{2} \bar{v}_{12,1'2'} \delta_{33'} \delta_{44'} \langle c_{1'} c_{2'} c_{3'} c_{4'} \rangle. \quad (18)$$

The graphical representation of the approximate $M^{\rm quartet}$ is shown in Fig. 4.

In Eq. (18), we put a renormalization factor λ' in front of the vertex which shall effectively account for the approximations considered, see App. D. It is, however, very fortunate that the final result is independent of the value of λ' and, therefore, one also may drop it. This somewhat surprising effect is due to selfconsistency and self-readjustment of the solution. It is demonstrated in the Appendix E for the simple case of ordinary pairing.

The complexity of the calculation still is further much reduced for the order parameter (14) with our mean field ansatz projected on zero total momentum, as already very successfully employed in [21]

$$\begin{aligned} \phi_{\vec{k}_1\vec{k}_2,\vec{k}_3\vec{k}_4} &= \varphi(|\vec{k}_1|)\varphi(|\vec{k}_2|)\varphi(|\vec{k}_3|)\varphi(|\vec{k}_4|) \\ &\quad \times (2\pi)^3 \delta(\vec{k}_1 + \vec{k}_2 + \vec{k}_3 + \vec{k}_4). \end{aligned} \quad (19)$$

It should be pointed out that this product ansatz with four identical $0S$ single particle wave functions is typical for a ground state configuration of the quartet. Excited configurations with wave functions of higher nodal structures may eventually be envisaged for other physical situations. We also would like to mention that the momentum conserving δ function induces strong correlations among the four particles and is, therefore, a rather non trivial variational wave function.

For the two-body interaction of $v_{\vec{k}_1\vec{k}_2,\vec{k}_3\vec{k}_4}$ in Eq. (15), we employ the same separable form as done already in our previous publication on the quartet critical temperature in Ref. [21]:

$$\begin{aligned} v_{\vec{k}_1\vec{k}_2,\vec{k}_3\vec{k}_4} &= \lambda w(\frac{\vec{k}_1 - \vec{k}_2}{2}) w(\frac{\vec{k}_3 - \vec{k}_4}{2}) \\ &\quad \times (2\pi)\delta^3(\vec{k}_1 + \vec{k}_2 - \vec{k}_3 - \vec{k}_4) \end{aligned} \quad (20)$$

with the form factor $w(\vec{k}) = w(|\vec{k}|) = e^{-k^2/b^2}$. An example for strength and range parameters are given in Sec. VI below.

With these simplifications, the mass operator (17) is independent of spin and isospin, and therefore it can be reduced to the following four-fold integral

$$\begin{aligned} & M^{\rm quartet}(k_1,\omega) \\ &= \frac{1}{(4\pi^2)^4} \int_0^\infty dK K^2 \int_{-1}^1 dt_1 \int_0^\infty dk k^2 \int_{-1}^1 dt \frac{\bar{f}(|\vec{K} - \vec{k}_1|)\bar{f}(|\frac{\vec{K}}{2} + \vec{k}|)\bar{f}(|\frac{\vec{K}}{2} - \vec{k}|) + f(|\vec{K} - \vec{k}_1|)f(|\frac{\vec{K}}{2} + \vec{k}|)f(|\frac{\vec{K}}{2} - \vec{k}|)}{\omega + \varepsilon_{\vec{K}-\vec{k}_1} + \varepsilon_{\frac{\vec{K}}{2}+\vec{k}} + \varepsilon_{\frac{\vec{K}}{2}+\vec{k}}} \\ &\times \ (w(|\vec{k}_1 - \frac{\vec{K}}{2}|))^2 (\varphi(|\frac{\vec{K}}{2} + \vec{k}|))^2 (\varphi(|\frac{\vec{K}}{2} - \vec{k}|))^2 \left[\int_0^\infty dk' k'^2 \int_{-1}^1 dt' w(k') \varphi(|\frac{\vec{K}}{2} + \vec{k}'|) \varphi(|\frac{\vec{K}}{2} - \vec{k}'|) \right]^2, \end{aligned} \quad (21)$$

with $t_1 = (\vec{K}\cdot\vec{k}_1)/(Kk_1)$, $t = (\vec{K}\cdot\vec{k})/(Kk)$, and $t' = (\vec{K}\cdot\vec{k}')/(Kk')$. Here we represented the Fermi distribution

function as $f_1 \to f(k_1) = f(\varepsilon_1)$. In this expression any strength factor of the vertices has been dropped, see our above argument and Appendix E. For the imaginary part of this expression an energy conserving delta function comes instead of the full denominator and then the 4D integral can be reduced to a 3D one. How this goes in detail is again explained in Appendix F. The real part of $M^{\rm quartet}$ is then obtained from the imaginary part via a dispersion integral:

$$\begin{aligned} &{\rm Re}M^{\rm quartet}(k,\omega+i\eta) \\ &= -\mathcal{P}\int_{-\infty}^{\infty}\frac{d\omega'}{\pi}\frac{{\rm Im}M^{\rm quartet}(k,\omega'+i\eta)}{\omega-\omega'} \end{aligned} \tag{22}$$

where $\mathcal{P}$ denotes the Cauchy principal value.

The occupation numbers are finally obtained from

$$\begin{aligned} \rho(k) &= -\int\frac{d\omega}{2\pi}2{\rm Im}G(k,\omega+i\eta)f(\omega) \\ &= \int\frac{d\omega}{2\pi}\frac{-2{\rm Im}M^{\rm quartet}(k,\omega+i\eta)}{(\omega-\varepsilon(\vec{k})-{\rm Re}M^{\rm quartet}(k,\omega+i\eta))^2+({\rm Im}M^{\rm quartet}(k,\omega+i\eta))^2}f(\omega). \end{aligned} \tag{23}$$

The equation for the order parameter (10) is formally not changed from Eqs. (4)-(7) of [21] but the occupation numbers are calculated selfconsistently with above equation. For completeness we again give the equations for the single particle wave function $\varphi(k)$

$$\mathcal{A}(k)\varphi(k)+3\mathcal{B}(k)+3\mathcal{C}(k)\varphi(k)=0, \tag{24}$$

where

$$\begin{aligned} \mathcal{A}(k) &= \int\frac{d^3k_2}{(2\pi)^3}\frac{d^3k_3}{(2\pi)^3}\frac{d^3k_4}{(2\pi)^3}\Big(\frac{k^2}{2m}+\frac{k_2^2}{2m}+\frac{k_3^2}{2m}+\frac{k_4^2}{2m}-4\mu\Big)\varphi^2(k_2)\varphi^2(k_3)\varphi^2(k_4)(2\pi)^3\delta(\vec{k}+\vec{k}_2+\vec{k}_3+\vec{k}_4) \\ &= \frac{1}{(2\pi)^4}\int dKK^2\int_{-1}^{1}dt\Big(\frac{k^2}{2m}+\frac{3}{2m}P^2-4\mu\Big)\varphi^2(P)\int dk'k'^2\int_{-1}^{1}dt'\varphi^2(k')\varphi^2(P'), \end{aligned} \tag{25}$$

$$\begin{aligned} \mathcal{B}(k) &= \int\frac{d^3k_2}{(2\pi)^3}\frac{d^3k_3}{(2\pi)^3}\frac{d^3k_4}{(2\pi)^3}\frac{d^3k_1'}{(2\pi)^3}\frac{d^3k_2'}{(2\pi)^3}(1-\rho(k)-\rho(k_2))\lambda w(\frac{\vec{k}-\vec{k}_2}{2})w(\frac{\vec{k}_1'-\vec{k}_2'}{2})(2\pi)^3\delta(\vec{k}+\vec{k}_2-\vec{k}_1'-\vec{k}_2') \\ &\quad\times\varphi(k_1')\varphi(k_2)\varphi(k_2')\varphi^2(k_3)\varphi^2(k_4)(2\pi)^3\delta(\vec{k}+\vec{k}_2+\vec{k}_3+\vec{k}_4) \\ &= \frac{\lambda}{(2\pi)^6}\int dKK^2\int_{-1}^{1}dt\Big(1-\rho(k)-\rho(P)\Big)w(Q)\varphi(P)\int dk'k'^2\int_{-1}^{1}dt'w(Q')\varphi(k')\varphi(P') \\ &\quad\times\int dk''k''^2\int_{-1}^{1}dt''\varphi^2(k'')\varphi^2(P''), \end{aligned} \tag{26}$$

$$\begin{aligned} \mathcal{C}(k) &= \int\frac{d^3k_2}{(2\pi)^3}\frac{d^3k_3}{(2\pi)^3}\frac{d^3k_4}{(2\pi)^3}\frac{d^3k_2'}{(2\pi)^3}\frac{d^3k_3'}{(2\pi)^3}(1-\rho(k_2)-\rho(k_3))\lambda w(\frac{\vec{k}_2-\vec{k}_3}{2})w(\frac{\vec{k}_2'-\vec{k}_3'}{2})(2\pi)^3\delta(\vec{k}_2+\vec{k}_3-\vec{k}_2'-\vec{k}_3') \\ &\quad\times\varphi(k_2)\varphi(k_2')\varphi(k_3)\varphi(k_3')\varphi^2(k_4)(2\pi)^3\delta(\vec{k}+\vec{k}_2+\vec{k}_3+\vec{k}_4) \\ &= \frac{\lambda}{(2\pi)^6}\int dKK^2\int_{-1}^{1}dt\varphi^2(P)\int dk'k'^2\int_{-1}^{1}dt'\Big(1-\rho(k')-\rho(P')\Big)w(Q')\varphi(k')\varphi(P') \\ &\quad\times\int dk''k''^2\int_{-1}^{1}dt''w(Q'')\varphi(k'')\varphi(P''). \end{aligned} \tag{27}$$

with

$$\begin{aligned} P &= \sqrt{K^2+k^2+2Kkt}, \\ P' &= \sqrt{K^2+k'^2+2Kk't'}, \end{aligned}$$

$$
\begin{aligned}
P'' &= \sqrt{K^2 + k''^2 + 2Kk''t''}, \\
Q &= \sqrt{K^2/4 + k^2 + Kkt}, \\
Q' &= \sqrt{K^2/4 + k'^2 + Kk't'}, \\
Q'' &= \sqrt{K^2/4 + k''^2 + Kk''t''}.
\end{aligned}
$$

As mentioned, in these equations the occupation numbers $\rho(k)$ shall be calculated selfconsistently from Eq. (23).

Because of its particular importance, before the presentation of the results, we first will discuss in the following section the 3h level density.

V. THREE HOLE LEVEL DENSITY

In what follows a crucial role will be played by the 3 hole propagator entering the mass operator. Since its influence on quartet condensation will be radically different from the corresponding 1h propagator in the pairing case, we will pay special attention to it in this section. In mean field approximation we can write

$$
\frac{(\bar{f}_1\bar{f}_2\bar{f}_3 + f_1f_2f_3)}{\omega + \varepsilon_{123}} = G^{(3h)}(k_1, k_2, k_3; \omega), \tag{28}
$$

where at r.h.s. we dropped spin and isospin indices since we consider unpolarised (nuclear) matter. We immediately see a strong difference with the pairing case. There, only a single hole line enters whose numerator is (see previous section) $\bar{f}_1 + f_1 = 1$, see Eq. (2) Therefore, no single particle occupation numbers appear in the numerator of a single hole propagator. This difference between the three hole and one hole propagators leads to strong consequences. This is best demonstrated in Fig. 5 with the three hole level density

$$
\begin{aligned}
& g(\omega) \\
&= -\int \frac{d^3k_1}{(2\pi)^3}\frac{d^3k_2}{(2\pi)^3}\frac{d^3k_3}{(2\pi)^3}\mathrm{Im}G^{(3h)}(k_1, k_2, k_3; \omega + i\eta) \\
&= \int \frac{d^3k_1}{(2\pi)^3}\frac{d^3k_2}{(2\pi)^3}\frac{d^3k_3}{(2\pi)^3} \\
&\quad \times(\bar{f}_1\bar{f}_2\bar{f}_3 + f_1f_2f_3)\pi\delta(\omega + \varepsilon_1 + \varepsilon_2 + \varepsilon_3) \\
&= \frac{m}{(2\pi)^5}\int_0^{k_{\max}} dkk^2 \int_0^{K_{\max}} dKK^2p \\
&\quad \times\Big(\bar{f}(k)\bar{F}(K,p) + f(k)F(K,p)\Big),
\end{aligned} \tag{29}
$$

where

$$
k_{\max} = \sqrt{2m(3\mu - \omega)}, \tag{30}
$$

$$
K_{\max} = \sqrt{4m(3\mu - \omega) - 2k^2}, \tag{31}
$$

$$
p = \sqrt{3m\mu - m\omega - \frac{k^2}{2} - \frac{K^2}{4}}, \tag{32}
$$



FIG. 5: 3h evel densities defined in Eq. (29) for various values of the chemical potential μ at a zero temperature.

and

$$
F(K,p) = \int_{-1}^{1} dtf(|\frac{\vec{K}}{2} + \vec{p}|)f(|\frac{\vec{K}}{2} - \vec{p}|), \tag{33}
$$

$$
\bar{F}(K,p) = \int_{-1}^{1} dt\bar{f}(|\frac{\vec{K}}{2} + \vec{p}|)\bar{f}(|\frac{\vec{K}}{2} - \vec{p}|) \tag{34}
$$

with $t = \vec{K}\cdot\vec{p}/(Kp)$.

In Fig. 5 we show the level density at zero temperature ($f(\omega) = \theta(-\omega)$), where it is calculated with the proton mass $m = 938.27$MeV (natural unit). Two cases have to be considered, chemical potential μ positive or negative. In the latter case we have binding of the quartet. Let us first discuss the case $\mu > 0$. We remark that in this case, the 3h level density goes through zero at $\omega = 0$, i.e. just in the region where the quartet correlations should appear. This is a strong difference with the pairing case where the 1h level density does not feel any influence from the medium and, therefore, the corresponding level density varies (neglecting the mean field for the sake of the argument) like in free space with the square root of energy. In particular, this means that the level density is *finite* at the Fermi level. This is a dramatic difference



FIG. 6: Single particle wave function $\varphi(k)$ in k-space (left), for r-space $\tilde{\varphi}(r)$ (middle), and occupation numbers (right) at $\mu = -5.26$ (top), -1.63 (middle) and 0.55 (bottom). The r-space wave function $\tilde{\varphi}(r)$ is derived from the Fourier transform of $\varphi(k)$ by $\tilde{\varphi}(r) = \int d^3k e^{i\vec{k}\cdot\vec{r}}\varphi(k)/(2\pi)^3$. The dashed line in the left panels correspond to the Gaussian with same norm and rms momentum as $\varphi(k)$.



FIG. 7: $-\mathrm{Im}M^{\mathrm{quartet}}(k_1, \omega + i\eta)$ in Eq. (21) as a function of ω for $\mu = -4.9$MeV (left) and for $\mu = 0.55$MeV (right) at zero temperature.

with the quartet case and explains why Cooper pairs can strongly overlap whereas for quartets this is impossible as we will see below. We also would like to point out that the 3h level density is just the mirror to the 3p level density which has been discussed in [28].

For the case where $\mu < 0$ there is nothing very special, besides the fact that it only is non-vanishing for negative values of ω and that the upper boundary is given by $\omega = 3\mu$. Therefore, the level density of Eq. (29) is zero for $\omega > 3\mu$.

VI. RESULTS AND DISCUSSION

At first let us mention that in this pilot application of our selfconsistent quartet theory, we only will consider the zero temperature case. As a definite physical example, we will treat the case of nuclear physics with the particularly strongly bound quartet, the α particle. It should be pointed out, however, that if scaled appropriately all energies and lengths can be transformed to other physical systems. For the nuclear case it is convenient to measure energies in in Fermi energies $\varepsilon_F = 35$MeV and lengths in inverse Fermi momentum $k_F^{-1} = 1.35^{-1}$fm.

To determine the order parameter, the calculation iterates in the following cycle:

1. Fix a chemical potential μ



FIG. 8: $-2\mathrm{Im}G(k, \omega + i\eta)$ in Eq. (23) as function of ω for $\mu = -4.9$MeV (top) and for $\mu = 0.55$MeV (bottom) at a zero temperature.

2. Give an initial trial order parameter $\varphi(k)$

3. Calculate the mass operator $M^{\rm quartet}(k,\omega)$ of Eq. (21) with $\varphi(k)$

4. Calculate the occupation numbers $\rho(k)$ with $M^{\rm quartet}(k,\omega)$ from Eq. (23)

5. Substituting $\varphi(k)$ and $\rho(k)$ into Eqs. (25), (26), and (27), derive the "new" single particle wave function $\varphi(k)$ from

$$\varphi(k) = \frac{-3\mathcal{B}(k)}{\mathcal{A}(k) + 3\mathcal{C}(k)}. \tag{35}$$

6. Quit the cycle once $\varphi(k)$ has converged, while go to 3. otherwise.

The single particle wave functions and occupation numbers obtained from the above cycle are shown in Fig. 6. We take $\lambda = -992$MeV fm^3 and $b = 1.43$fm^{-1} to get the binding energy of the free α-particle (-28.3MeV) and its radius (1.7fm). We also insert the Gaussian wave function with same rms momentum as the single particle wave function in the left figures in Fig. 6. As shown in Fig. 6, the single particle wave function is sharper than a Gaussian. There is the term $\mathcal{A}(k)$ of Eq. (25) in the denominator of $\varphi(k)$ in Eq. (35), and $\mathcal{A}(k)$ involves the factor $(k^2/(2m) - 4\mu)$. Hence, $\varphi(k)$ is closer to a Lorentzian rather than to a Gaussian [1].

We could not obtain a convergent wave function for $\mu > 0.55$MeV. This difficulty is of the same origin as in the case of our calculation of the critical temperature for α particle condensation. This stems from the fact that for larger positive values of the chemical potential, the denominator of $\varphi(k)$ in Eq. (35) at a certain value of k becomes zero, while the numerator is finite. In case of pair condensation, as shown in Eq. (4), both the denominator and numerator become zero at the same value of k, this being a further crucial difference between the quartet and pairing cases. In the r.h.s. panels of Fig. 6 we also show the corresponding occupation numbers. We see that they are very small. However, they increase for increasing values of the chemical potential. For $\mu = 0.55$ MeV the maximum of the occupation still only attains 0.35 what is far away from the saturation value of one. What really happens for larger values of the chemical potential, is unclear. Surely, as discussed in Sec. V the situation for the quartet case is completely different from the standard pairing case. This is due to the fact, as already mentioned, that the 3h level density goes through zero at $\omega = 0$, i.e. just at the place where the quartet correlation should build up for positive values of μ. Due to this fact, the inhibition to go into the positive μ regime is here even stronger than in the case of the critical temperature [21].

The situation in the quartet case is also in so far much different, as the the 3h GF produces a considerable imaginary part of the mass operator. Figure 7 shows the imaginary part of the approximate quartet mass operator of Eq. (21) for $\mu < 0$ and $\mu > 0$. These large values of the damping rate imply a strong violation of the quasiparticle picture. In Fig. 8 we show the spectral function of the single particle GF. Contrary to the pairing case with its sharp quasiparticle pole, we here only find a very broad distribution, implying that the quasiparticle picture is completely destroyed. How to formulate a theory which goes continuously from the quartet case into the pairing case, is an open question. One solution could be to start right from the beginning with an in medium four body equation which contains a superfluid phase. When the quartet phase disappears, the superfluid phase may remain. Such investigations shall be done in the future.

VII. CONCLUSION

We formulated the gap equation for quartetting in fermion systems in analogy to the BCS gap equation. The mass operator of quartet-BEC with effective four-body vertices and in-medium four-body Schrödinger equation was derived with the Dyson equation approach to correlation functions. However, the full expression of the quartet mass operator is too complicated to be evaluated numerically in a direct manner. The biggest problem stems from the many dimensional integrals over momenta. We, therefore, introduced some reasonable approximations reducing the complexity considerably. A feature of great help is that the final answer is independent of the strength of the vertex function, making our approximation probably quite reliable. In our calculation we also applied the same mean field ansatz projected on zero total momentum which was already so successful in our previous calculation of the critical temperature [21]. This feature, of course, reduces the numerical effort tremendously, since only a single 0S wave function has to be determined selfconsistently by iteration.

In this pilot work with an application to nuclear physics, we showed results only at zero temperature, however, the formalism we presented is at finite temperature.

We think that the results are of general validity showing qualitatively very distinct features from the pairing case. For example no well defined quasiparticles occur in the case of quartets. This is due to the fact that the quartet order parameter in the single particle mass operator goes along with the level density of three uncorrelated holes. Only the total momentum of the three holes is well defined and equal to the time reversed momentum of the incoming particle. The relative momenta of the three holes have to be integrated over, yielding a strong imaginary part of the mass operator smoothing out any individual single particle structure. Another remarkable feature already encountered in our previous work in Ref. [21] is that the self consistent solution seems to exist only from negative chemical potential until around zero, i.e. from the BEC, or strong coupling region until crossover region. Once one goes to positive μ's the solution breaks down. This effect is even more pronounced here than it

was in our study [21] for the critical temperature. This can be traced back to the fact that the 3h level density goes through zero at 3μ for $\mu > 0$, that is just at the place where the quartet correlations should occur. Actually this feature is present for all multi-particle multi-hole level densities. The only exception being the single particle level density which is finite at the Fermi level. This unique feature makes that pairing also is unique and for instance allows for a weak coupling situation with a coherence length of the pair orders of magnitude larger than the interparticle distance. It remains an open problem, how to formulate a more general theory which continuously goes from the quartet case to the pairing case.

Acknowledgments

P. S. wants to thank M. Urban for useful discussions. This work is supported by the DFG Grant No. RO905/29-1.

Appendix A: Dyson equations for multi-particle-hole Green functions

We shall shortly review the basic formulation used in the present work. We here extend the approach for real-time Green's functions (GF's) at zero temperature in Ref. [30] to real-time GF's at finite temperature [25, 31].

The Hamiltonian in a fermion system with two-body interaction is

$$\begin{aligned} & K = H - \mu N = T + V - \mu N \\ = & \sum_1 \varepsilon_1 c_1^\dagger c_1 + \frac{1}{4} \sum_{1234} \bar{v}_{12,34} c_1^\dagger c_2^\dagger c_4 c_3. \end{aligned} \tag{A1}$$

The c_1, $c_1^\dagger$ are fermion annihilation and creation operators with an arbitrary quantum number 1.

A real-time multi-particle-hole GF at a finite temperature is defined by

$$\begin{aligned} & G^{(i\mathrm{p}j\mathrm{h})}_{\alpha;\alpha'}(t-t') \\ = & \begin{cases} -i\langle T(A_\alpha(t) A^\dagger_{\alpha'}(t'))\rangle & \text{(chronological)} \\ -i\theta(t-t')\langle [A_\alpha(t), A^\dagger_{\alpha'}(t')]_\pm\rangle & \text{(retarded)} \end{cases} \end{aligned} \tag{A2}$$

where $\langle ...\rangle$ means the thermal average, T is the time ordering operator, A_α is an arbitrary operator consisting out of individual fermion operators c_1 and $c_1^\dagger$, and $[...,...]_\pm$ is the anti-commutator or commutator. The time dependence of the operators is given in the Heisenberg picture $A_\alpha(t) = e^{iKt} A_\alpha e^{-iKt}$.

Note that, although we treat chronological GF's below, the change to retarded, advanced and Matsubara GF's goes as usual [25]. Below, when we go from time space to Fourier space, we always will go over to retarded GF's without mentioning it explicitly.

The superscript $(i\mathrm{p}j\mathrm{h})$ in Eq. (A2) means i-particle j-hole GF, where i (j) is the number of the annihilation (creation) operators in A_α, e.g. $G^{(1\mathrm{p})}_{1;1'}(t-t') = -i\langle T(c_1(t) c_{1'}^\dagger(t'))\rangle$, $G^{(1\mathrm{p}1\mathrm{h})}_{1,2;1',2'}(t-t') = -i\langle T((c_2^\dagger c_1)_t (c_{1'}^\dagger c_{2'})_{t'})\rangle$ with $(c_2^\dagger c_1)_t = c_2^\dagger(t) c_1(t)$, etc..

The Dyson equation for the $i\mathrm{p}j\mathrm{h}$ GF is [30]

$$\begin{aligned} & (i\frac{\partial}{\partial t} - \varepsilon_\alpha) G^{(i\mathrm{p}j\mathrm{h})}_{\alpha;\alpha'}(t-t') \\ = & \ \delta(t-t')\mathcal{N}_{\alpha;\alpha'} + \sum_\beta M^0_{\alpha;\beta} G^{(i\mathrm{p}j\mathrm{h})}_{\beta;\alpha'}(t-t') \\ + & \sum_\beta \int dt'' M_{\alpha;\beta}(t-t'') G^{(i\mathrm{p}j\mathrm{h})}_{\beta;\alpha'}(t''-t'), \end{aligned} \tag{A3}$$

with

$$M^0_{\alpha;\alpha'} = \sum_\beta \langle [[A_\alpha, V]_-, A^\dagger_\beta]_\pm\rangle \mathcal{N}^{-1}_{\beta;\alpha'} \tag{A4}$$

$$\begin{aligned} & M_{\alpha;\alpha'}(t-t') \\ = & -i\sum_\beta \langle T([A_\alpha(t), V]_- [V, A^\dagger_\beta(t')]_-)\rangle_{\mathrm{irr.}} \mathcal{N}^{-1}_{\beta;\alpha'}, \end{aligned} \tag{A5}$$

$$\mathcal{N}_{\alpha;\alpha'} = \langle [A_\alpha, A^\dagger_{\alpha'}]_\pm\rangle \tag{A6}$$

where ε_α is defined by $[A_\alpha, T-\mu N]_- = \varepsilon_\alpha A_\alpha$, and $\mathcal{N}^{-1}_{\alpha;\alpha'}$ is the inverse of the matrix $\mathcal{N}_{\alpha;\alpha'}$.

According to the time dependence, we shall call $M^0_{\alpha;\alpha'}$ static mass operator and $M_{\alpha;\alpha'}(t-t')$ dynamical mass operator.

Since $[A_\alpha(t), V]_-$ is the operator A_α augmented by one annihilation and one creation operator, $\langle T([A_\alpha(t), V]_- [V, A^\dagger_\beta(t'')]_-)\rangle_{\mathrm{irr.}}$ becomes a $(i{+}1)\mathrm{p}(j{+}1)\mathrm{h}$ GF in the dynamical mass of Eq. (A5). The index 'irr.' in Eq. (A5) stands for the mass operator being irreducible with respect to a cut of $i\mathrm{p}j\mathrm{h}$ lines [26].

From the Dyson equation of Eq. (A3) we can see that the Fourier transform of the bare $i\mathrm{p}j\mathrm{h}$ GF is

$$G^{0(i\mathrm{p}j\mathrm{h})}_{\alpha;\alpha'}(\omega) = \frac{\mathcal{N}_{\alpha;\alpha'}}{\omega - \varepsilon_\alpha}. \tag{A7}$$

Appendix B: Quartet mass operator

Here we derive the quartet mass operator of Eq. (8) shown in Sec. III.

Notice that we shall use summation convention for repeated indices and neglect all terms except the ones which are associated with the quartet order parameters.

From Eq. (A3), the Dyson equation for the 1p GF is [26]

$$\begin{aligned} & \left(i\frac{\partial}{\partial t} - \varepsilon_1\right) G^{(1\mathrm{p})}_{1;2}(t_1 - t_2) \\ = & \ \delta_{12}\delta(t_1 - t_2) \\ + & \int dt_3 M^{(1\mathrm{p})}_{1;3}(t_1 - t_3) G^{(1\mathrm{p})}_{3;2}(t_3 - t_2). \end{aligned} \tag{B1}$$

The Fourier transform yields

$$(\omega-\varepsilon_1)G^{(1\mathrm{p})}_{1;2}(\omega)=\delta_{12}+M^{(1\mathrm{p})}_{1;3}(\omega)G^{(1\mathrm{p})}_{3;2}(\omega) \quad \text{(B2)}$$

with

$$M^{(1\mathrm{p})}_{1;2}(\omega)=\frac{1}{2}\bar{v}_{1z_1,a_1a_1'}G^{(2\mathrm{p1h})\mathrm{irr.}}_{a_1a_1',z_1;a_2a_2',z_2}(\omega)\frac{1}{2}\bar{v}_{2z_2,a_2a_2'}. \quad \text{(B3)}$$

Therefore

$$\begin{aligned}&G^{(1\mathrm{p})}_{1;2}(\omega)\\ =\;& G^{0(1\mathrm{p})}_{1;2}(\omega)+G^{0(1\mathrm{p})}_{1;3}(\omega)M^{(1\mathrm{p})}_{3;4}(\omega)G^{(1\mathrm{p})}_{4;2}(\omega), \quad \text{(B4)}\end{aligned}$$

where

$$G^{0(1\mathrm{p})}_{1;2}(\omega)=\frac{\delta_{12}}{\omega-\varepsilon_1}. \quad \text{(B5)}$$

For the Dyson equation of the 2p1h GF one obtains

$$\begin{aligned}&\left(\omega-(\varepsilon_{a_1}+\varepsilon_{a_1'}-\varepsilon_{z_1})\right)G^{(2\mathrm{p1h})}_{a_1a_1',z_1;a_2a_2',z_2}(\omega)\\ =\;&\mathcal{N}_{a_1a_1',z_1;a_2a_2',z_2}\\ +\;&M^{(2\mathrm{p1h})}_{a_1a_1',z_1;a_3a_3',z_3}(\omega)G^{(2\mathrm{p1h})}_{a_3a_3',z_3;a_2a_2',z_2}(\omega), \quad \text{(B6)}\end{aligned}$$

where

$$\begin{aligned}&M^{(2\mathrm{p1h})}_{a_1a_1',z_1;a_2a_2',z_2}(\omega)\\ =\;&\Big[\frac{1}{2}\bar{v}_{a_1y_1,b_1b_1'}\delta_{z_1y_1'}\delta_{a_1'b_1''}+\frac{1}{2}\bar{v}_{a_1'y_1',b_1b_1'}\delta_{z_1y_1}\delta_{a_1b_1''}\\ &-\frac{1}{2}\bar{v}_{z_1b_1'',y_1y_1'}\delta_{a_1b_1}\delta_{a_1'b_1'}\Big]G^{(3\mathrm{p2h})\mathrm{irr.}}_{b_1b_1'b_1'',y_1y_1';b_2b_2'b_2'',y_2y_2'}(\omega)\\ \times\;&\Big[\frac{1}{2}\bar{v}_{a_3y_2,b_2b_2'}\delta_{z_3y_2'}\delta_{a_3'b_2''}+\frac{1}{2}\bar{v}_{a_3'y_2',b_2b_2'}\delta_{z_3y_2}\delta_{a_3b_3''}\\ &-\frac{1}{2}\bar{v}_{z_3b_2'',y_2y_2'}\delta_{a_3b_2}\delta_{a_3'b_2'}\Big]\mathcal{N}^{-1}_{a_3a_3',z_3;a_2a_2',z_2} \quad \text{(B7)}\end{aligned}$$

and

$$\mathcal{N}_{a_1a_1',z_1;a_2a_2',z_2}=\langle[c^\dagger_{z_1}c_{a_1'}c_{a_1},c^\dagger_{a_2}c^\dagger_{a_2'}c_{z_2}]_+\rangle. \quad \text{(B8)}$$

Therefore

$$\begin{aligned}&G^{(2\mathrm{p1h})}_{a_1a_1',z_1;a_2a_2',z_2}(\omega)\\ =\;&G^{0(2\mathrm{p1h})}_{a_1a_1',z_1;a_2a_2',z_2}(\omega)\\ +\;&G^{0(2\mathrm{p1h})}_{a_1a_1',z_1;a_3a_3',z_3}(\omega)\mathcal{N}^{-1}_{a_3a_3',z_3;a_4a_4',z_4}\\ &\times M^{(2\mathrm{p1h})}_{a_4a_4',z_4;a_5a_5',z_5}(\omega)G^{(2\mathrm{p1h})}_{a_5a_5',z_5;a_2a_2',z_2}(\omega), \quad \text{(B9)}\end{aligned}$$

with

$$G^{0(2\mathrm{p1h})}_{a_1a_1',z_1;a_2a_2',z_2}(\omega)=\frac{\mathcal{N}_{a_1a_1',z_1;a_2a_2',z_2}}{\omega-(\varepsilon_{a_1a_1'}-\varepsilon_{z_1})} \quad \text{(B10)}$$

and

$$\mathcal{N}_{a_1a_1',z_1;a_3a_3',z_3}\mathcal{N}^{-1}_{a_3a_3',z_3;a_2a_2',z_2}=\delta_{a_1a_2}\delta_{a_1'a_2'}\delta_{z_1z_2}, \quad \text{(B11)}$$



FIG. 9: Graphical representation of $\Gamma^{(3)}$ in Eq. (B13). The dots is represent the two-body interaction.

where $\varepsilon_{ij\ldots}=\varepsilon_i+\varepsilon_j+\cdots$ in (B10).

Substituting the 2p1h GF of Eq. (B9) into the mass operator of Eq. (B3) we obtain

$$\begin{aligned}&M^{(1\mathrm{p})}_{1;2}(\omega)\\ =\;&\Gamma^{(3)}_{1y_1y_1',b_1b_1'b_1''}(\omega)G^{(3\mathrm{p2h})\mathrm{irr.}}_{b_1b_1'b_1'',y_1y_1';b_2b_2'b_2'',y_2y_2'}(\omega)\\ &\times\Gamma^{(3)*}_{2y_2y_2',b_2b_2'b_2''}(\omega), \quad \text{(B12)}\end{aligned}$$

where

$$\begin{aligned}&\Gamma^{(3)}_{1y_1y_1',b_1b_1'b_1''}(\omega)\\ =\;&\frac{1}{2}\bar{v}_{1z_1,a_1a_1'}G^{0(2\mathrm{p1h})}_{a_1a_1',z_1;a_3a_3',z_3}(\omega)\mathcal{N}^{-1}_{a_3a_3',z_3;a_4a_4',z_4}\\ \times\;&\Big[\frac{1}{2}\bar{v}_{a_4y_1,b_1b_1'}\delta_{z_1y_1'}\delta_{a_4'b_1''}+\frac{1}{2}\bar{v}_{a_4'y_1',b_1b_1'}\delta_{z_1y_1}\delta_{a_4b_1''}\\ &-\frac{1}{2}\bar{v}_{z_4b_1'',y_1y_1'}\delta_{a_4b_1}\delta_{a_4'b_1'}\Big]. \quad \text{(B13)}\end{aligned}$$

In (B12) we omitted the term derived from the first term at r.h.s. of (B9) because it is disconnected with the quartet order parameter. In the effective three body vertex $\Gamma^{(3)}$ of Eq. (B13) we give not the exact 2p1h GF but the free one, as we only consider in this work the lowest order approximation, though the exact 2p1h GF figures, in principle, in the right vertex $\Gamma^{(3)}$ of Eq. (B13) substituting (B9) into (B3). In Eq. (B13), using Eqs. (B10) and (B11), we obtain

$$\begin{aligned}&G^{0(2\mathrm{p1h})}_{a_1a_1',z_1;a_3a_3',z_3}(\omega)\mathcal{N}^{-1}_{a_3a_3',z_3;a_4a_4',z_4}\\ =\;&\frac{\delta_{a_1a_4}\delta_{a_1'a_4'}\delta_{z_1z_4}}{\omega-\varepsilon_{a_1a_1'}+\varepsilon_{z_1}}, \quad \text{(B14)}\end{aligned}$$

and thus Eq. (B13) is consistent with Eq. (11).

Furthermore, the Dyson equation for 3p2h GF is

$$\begin{aligned}&(\omega-(\varepsilon_{b_1b_1'b_1''}-\varepsilon_{y_1y_1'}))G^{(3\mathrm{p2h})}_{b_1b_1'b_1'',y_1y_1';b_2b_2'b_2'',y_2y_2'}(\omega)\\ =\;&\mathcal{N}_{b_1b_1'b_1'',y_1y_1';b_2b_2'b_2'',y_2y_2'}\\ +\;&M^{(3\mathrm{p2h})}_{b_1b_1'b_1'',y_1y_1';b_3b_3'b_3'',y_3y_3'}(\omega)\\ &\times G^{(3\mathrm{p2h})}_{b_3b_3'b_3'',y_3y_3';b_2b_2'b_2'',y_2y_2'}(\omega). \quad \text{(B15)}\end{aligned}$$

The mass operator is then given by

$$M^{(3\mathrm{p2h})}_{b_1b_1'b_1'',y_1y_1';b_2b_2'b_2'',y_2y_2'}(\omega)$$

$$
\begin{aligned}
&= \Big[\frac{1}{2}\bar{v}_{b_1x_1'',c_1c_1'''}\delta_{y_1x_1}\delta_{y_1'x_1'}\delta_{b_1'c_1'}\delta_{b_1''c_1''} \\
&\quad+\frac{1}{2}\bar{v}_{b_1'x_1'',c_1'c_1'''}\delta_{y_1x_1}\delta_{y_1'x_1'}\delta_{b_1c_1}\delta_{b_1''c_1''} \\
&\quad+\frac{1}{2}\bar{v}_{b_1''x_1'',c_1''c_1'''}\delta_{y_1x_1}\delta_{y_1'x_1'}\delta_{b_1c_1}\delta_{b_1'c_1'} \\
&\quad-\frac{1}{2}\bar{v}_{y_1c_1''',x_1x_1''}\delta_{y_1'x_1'}\delta_{b_1c_1}\delta_{b_1'c_1'}\delta_{b_1''c_1''} \\
&\quad-\frac{1}{2}\bar{v}_{y_1'c_1''',x_1'x_1''}\delta_{y_1x_1}\delta_{b_1c_1}\delta_{b_1'c_1'}\delta_{b_1''c_1''}\Big] \\
&\quad\times G^{(4\mathrm{p3h})\mathrm{irr.}}_{c_1c_1'c_1''c_1''',x_1x_1'x_1'';c_2c_2'c_2''c_2''',x_2x_2'x_2''}(\omega) \\
&\times \Big[\frac{1}{2}\bar{v}_{b_3x_2'',c_2c_2'''}\delta_{y_3x_2}\delta_{y_3'x_2'}\delta_{b_3'c_2'}\delta_{b_3''c_2''} \\
&\quad+\frac{1}{2}\bar{v}_{b_3'x_2'',c_2'c_2'''}\delta_{y_3x_2}\delta_{y_3'x_2'}\delta_{b_3c_2}\delta_{b_3''c_2''} \\
&\quad+\frac{1}{2}\bar{v}_{b_3''x_2'',c_2''c_2'''}\delta_{y_3x_2}\delta_{y_3'x_2'}\delta_{b_3c_2}\delta_{b_3'c_2'} \\
&\quad-\frac{1}{2}\bar{v}_{y_3c_2''',x_2x_2''}\delta_{y_3'x_2'}\delta_{b_3c_2}\delta_{b_3'c_2'}\delta_{b_3''c_2''} \\
&\quad-\frac{1}{2}\bar{v}_{y_3'c_2''',x_2'x_2''}\delta_{y_3x_2}\delta_{b_3c_2}\delta_{b_3'c_2'}\delta_{b_3''c_2''}\Big] \\
&\quad\times\mathcal{N}^{-1}_{b_3b_3'b_3'',y_3y_3';b_2b_2'b_2'',y_2y_2'}, \qquad \text{(B16)}
\end{aligned}
$$

and

$$
\begin{aligned}
&\mathcal{N}_{b_1b_1'b_1'',y_1y_1';b_2b_2'b_2'',y_2y_2'} \\
&= \langle[c^\dagger_{y_1}c^\dagger_{y_1'}c_{b_1''}c_{b_1'}c_{b_1},c^\dagger_{b_1}c^\dagger_{b_1'}c^\dagger_{b_1''}c_{y_1'}c_{y_1}]_+\rangle. \qquad \text{(B17)}
\end{aligned}
$$

Therefore, one obtains for the 3p2h GF

$$
\begin{aligned}
&G^{(3\mathrm{p2h})}_{b_1b_1'b_1'',y_1y_1';b_2b_2'b_2'',y_2y_2'}(\omega) \\
&= G^{0(3\mathrm{p2h})}_{b_1b_1'b_1'',y_1y_1';b_2b_2'b_2'',y_2y_2'}(\omega) \\
&+ G^{0(3\mathrm{p2h})}_{b_1b_1'b_1'',y_1y_1';b_3b_3'b_3'',y_3y_3'}(\omega)\mathcal{N}^{-1}_{b_3b_3'b_3'',y_3y_3';b_4b_4'b_4'',y_4y_4'} \\
&\quad\times M^{(3\mathrm{p2h})}_{b_4b_4'b_4'',y_4y_4';b_5b_5'b_5'',y_5y_5'}(\omega) \\
&\quad\times G^{(3\mathrm{p2h})}_{b_5b_5'b_5'',y_5y_5';b_2b_2'b_2'',y_2y_2'}(\omega), \qquad \text{(B18)}
\end{aligned}
$$

where

$$
\begin{aligned}
&G^{0(3\mathrm{p2h})}_{b_1b_1'b_1'',y_1y_1';b_2b_2'b_2'',y_2y_2'}(\omega) \\
&= \frac{\mathcal{N}_{b_1b_1'b_1'',y_1y_1';b_2b_2'b_2'',y_2y_2'}}{\omega-(\varepsilon_{b_1b_1'b_1''}-\varepsilon_{y_1y_1'})}. \qquad \text{(B19)}
\end{aligned}
$$

Substituting 3p2h GF of (B18) into the mass operator of Eq. (B12) leads to

$$
\begin{aligned}
&M^{(1\mathrm{p})}_{1;2}(\omega) \\
&= \Gamma^{(4)}_{1x_1x_1'x_1'',c_1c_1'c_1''c_1'''}(\omega) \\
&\quad\times G^{(4\mathrm{p3h})\mathrm{irr.}}_{c_1c_1'c_1''c_1''',x_1x_1'x_1'';c_2c_2'c_2''c_2''',x_2x_2'x_2''}(\omega) \\
&\quad\times\Gamma^{(4)*}_{2x_2x_2'x_2'',c_2c_2'c_2''c_2'''}(\omega), \qquad \text{(B20)}
\end{aligned}
$$



FIG. 10: Graphical representation of $\Gamma^{(4)}$ in Eq. (B21).

with

$$
\begin{aligned}
&\Gamma^{(4)}_{1x_1x_1'x_1'',c_1c_1'c_1''c_1'''}(\omega) \\
&= \Gamma^{(3)}_{1y_1y_1',b_1b_1'b_1''}(\omega) \\
&\quad\times G^{0(3\mathrm{p2h})}_{b_1b_1'b_1'',y_1y_1';b_3b_3'b_3'',y_3y_3'}(\omega)\mathcal{N}^{-1}_{b_3b_3'b_3'',y_3y_3';b_4b_4'b_4'',y_4y_4'} \\
&\times \Big[\frac{1}{2}\bar{v}_{b_4x_1'',c_1c_1'''}\delta_{y_4x_1}\delta_{y_4'x_1'}\delta_{b_4'c_1'}\delta_{b_4''c_1''} \\
&\quad+\frac{1}{2}\bar{v}_{b_4'x_1'',c_1'c_1'''}\delta_{y_4x_1}\delta_{y_4'x_1'}\delta_{b_4c_1}\delta_{b_4''c_1''} \\
&\quad+\frac{1}{2}\bar{v}_{b_4''x_1'',c_1''c_1'''}\delta_{y_4x_1}\delta_{y_4'x_1'}\delta_{b_4c_1}\delta_{b_4'c_1'} \\
&\quad-\frac{1}{2}\bar{v}_{y_4c_1''',x_1x_1''}\delta_{y_4'x_1'}\delta_{b_4c_1}\delta_{b_4'c_1'}\delta_{b_4''c_1''} \\
&\quad-\frac{1}{2}\bar{v}_{y_4'c_1''',x_1'x_1''}\delta_{y_4x_1}\delta_{b_4c_1}\delta_{b_4'c_1'}\delta_{b_4''c_1''}\Big] \qquad \text{(B21)}
\end{aligned}
$$

Here we also omitted the terms associated with the first term at r.h.s. of Eq. (B18) because it is disconnected with the quartet order parameter. Besides we introduced the free 3p2h GF in $\Gamma^{(4)}$ of (B21) for the same reason as this was done in Eqs. (B12) and (B13). The graphical representation of $\Gamma^{(4)}$ is shown in Fig. 10.

The 4p3h GF can, for our purpose of α-particle condensation, approximately be decomposed into the order parameter and the free three body GF as follows (in analogy to what is done in the pairing case)

$$
\begin{aligned}
&G^{(4\mathrm{p3h})}_{c_1c_1'c_1''c_1''',x_1x_1'x_1'';c_2c_2'c_2''c_2''',x_2x_2'x_2''}(t_1-t_2) \\
&= -i\langle T(c^\dagger_{x_1}c^\dagger_{x_1'}c^\dagger_{x_1''}c_{c_1'''}c_{c_1''}c_{c_1'}c_{c_1})_{t_1}(c^\dagger_{c_2}c^\dagger_{c_2'}c^\dagger_{c_2''}c^\dagger_{c_2'''}c_{x_2''}c_{x_2'}c_{x_2})_{t_2}\rangle \\
&= -i\langle T(c_{c_1'''}c_{c_1''}c_{c_1'}c_{c_1})_{t_1}\rangle\langle T(c^\dagger_{c_2}c^\dagger_{c_2'}c^\dagger_{c_2''}c^\dagger_{c_2'''})_{t_2}\rangle \\
&\quad\times\langle T(c^\dagger_{x_1}c^\dagger_{x_1'}c^\dagger_{x_1''})_{t_1}(c_{x_2''}c_{x_2'}c_{x_2})_{t_2}\rangle \\
&= \langle c_{c_1'''}c_{c_1''}c_{c_1'}c_{c_1}\rangle\langle c^\dagger_{c_2}c^\dagger_{c_2'}c^\dagger_{c_2''}c^\dagger_{c_2'''}\rangle \\
&\quad\times G^{0(3\mathrm{h})}_{x_1x_1'x_1'';x_2x_2'x_2''}(t_1-t_2) \qquad \text{(B22)}
\end{aligned}
$$

The Fourier transform is

$$
G^{(4\mathrm{p3h})}_{c_1c_1'c_1''c_1''',x_1x_1'x_1'';c_2c_2'c_2''c_2''',x_2x_2'x_2''}(\omega)
$$

$$= \langle c_{c_1'''} c_{c_1''} c_{c_1'} c_{c_1} \rangle \langle c^\dagger_{c_2} c^\dagger_{c_2'} c^\dagger_{c_2''} c^\dagger_{c_2'''} \rangle \times G^{0(3h)}_{x_1 x_1' x_1''; x_2 x_2' x_2''}(\omega). \quad \text{(B23)}$$

The free 3h GF is explicitly given by

$$G^{0(3h)}_{x_1 x_1' x_1''; x_2 x_2' x_2''}(\omega) = \frac{\bar{f}_{x_1} \bar{f}_{x_1'} \bar{f}_{x_1''} + f_{x_1} f_{x_1'} f_{x_1''}}{\omega + \varepsilon_{x_1 x_1' x_1''}} P_{x_1 x_1' x_1''; x_2 x_2' x_2''} \quad \text{(B24)}$$

with $P_{x_1 x_1' x_1''; x_2 x_2' x_2''}$ as in Eq. (9).

Therefore, Eq. (B23) becomes

$$G^{(4p3h)}_{c_1 c_1' c_1'' c_1''', x_1 x_1' x_1''; c_2 c_2' c_2'' c_2''', x_2 x_2' x_2''}(\omega) = \langle c_{c_1'''} c_{c_1''} c_{c_1'} c_{c_1} \rangle \langle c^\dagger_{c_2} c^\dagger_{c_2'} c^\dagger_{c_2''} c^\dagger_{c_2'''} \rangle \times \frac{\bar{f}_{x_1} \bar{f}_{x_1'} \bar{f}_{x_1''} + f_{x_1} f_{x_1'} f_{x_1''}}{\omega + \varepsilon_{x_1 x_1' x_1''}} P_{x_1 x_1' x_1''; x_2 x_2' x_2''}. \quad \text{(B25)}$$

Finally we obtain for the quartet mass operator,

$$M^{\text{quartet}}_{1;2}(\omega) = \Gamma^{(4)}_{1 x_1 x_1' x_1''; c_1 c_1' c_1'' c_1'''}(\omega) \langle c_{c_1'''} c_{c_1''} c_{c_1'} c_{c_1} \rangle \times \frac{\bar{f}_{x_1} \bar{f}_{x_1'} \bar{f}_{x_1''} + f_{x_1} f_{x_1'} f_{x_1''}}{\omega + \varepsilon_{x_1 x_1' x_1''}} P_{x_1 x_1' x_1''; x_2 x_2' x_2''} \times \langle c^\dagger_{c_2} c^\dagger_{c_2'} c^\dagger_{c_2''} c^\dagger_{c_2'''} \rangle \Gamma^{(4)*}_{2 x_2 x_2' x_2''; c_2 c_2' c_2'' c_2'''}(\omega). \quad \text{(B26)}$$

Appendix C: in-medium quartet order parameter equation

We here give a short derivation of the in-medium quartet order parameter equation. It is derived from the Dyson equation for 4p GF in static approximation of the mass operator. The Dyson equation for 4p GF derived from Eq. (A3) without dynamical mass operator is

$$\left(i\frac{\partial}{\partial t} - \varepsilon_{1234} \right) G^{(4p)}_{1234;1'2'3'4'}(t - t') = \mathcal{N}_{1234;1'2'3'4'} + V_{1234;1''2''3''4''} G^{(4p)}_{1234;1'2'3'4'}(t - t') \quad \text{(C1)}$$

with in the ladder approximation

$$\begin{aligned} & V_{1234;1'2'3'4'} \\ &= \langle [[c_4 c_3 c_2 c_1, V]_-, c^\dagger_{1''} c^\dagger_{2''} c^\dagger_{3''} c^\dagger_{4''}]_- \rangle \mathcal{N}^{-1}_{1''2''3''4'';1'2'3'4'} \\ &= \frac{1}{2}(1 - \rho_1 - \rho_2) \bar{v}_{12,1'2'} \delta_{33'} \delta_{44'} \\ &+ \frac{1}{2}(1 - \rho_1 - \rho_3) \bar{v}_{13,1'3'} \delta_{22'} \delta_{44'} \\ &+ \frac{1}{2}(1 - \rho_1 - \rho_4) \bar{v}_{14,1'4'} \delta_{22'} \delta_{33'} \\ &+ \text{permutations} \end{aligned} \quad \text{(C2)}$$

where we approximated the correlation functions by factorizing them into products of single particle occupation numbers.

The Fourier transform of 4p GF is

$$\begin{aligned} & G^{(4p)}_{1234;1'2'3'4'}(\omega) \\ &= G^{0(4p)}_{1234;1'2'3'4'}(\omega) \\ &+ G^{0(4p)}_{1234;1''2''3''4''}(\omega) \mathcal{N}^{-1}_{1''2''3''4'';5678} \\ &\quad \times V_{5678;5'6'7'8'} G^{(4p)}_{5'6'7'8';1'2'3'4'}(\omega), \end{aligned} \quad \text{(C3)}$$

where

$$G^{0(4p)}_{1234;1'2'3'4'}(\omega) = \frac{\mathcal{N}_{1234;1'2'3'4'}}{\omega - \varepsilon_{1234}}. \quad \text{(C4)}$$

From the spectral representation of the 4p GF we shall only retain the ground state because of its condensate character. Therefore, at the ground state pole, i.e. at $\omega = 0$, we obtain

$$\langle c_4 c_3 c_2 c_1 \rangle = -\frac{1}{\varepsilon_{1234}} V_{1234;1'2'3'4'} \langle c_{4'} c_{3'} c_{2'} c_{1'} \rangle. \quad \text{(C5)}$$

This equation corresponds to Eq. (4) in the pairing case.

Appendix D: Approximate mass operator in Eq. (17)

After the derivation of a single particle mass operator containing the quartet condensate in App. B, we easily recognize that its expression, for instance the vertex $\Gamma^{(4)}$ in Eqs. (10) and (11) (or (B13) and (B21)) it contains, is of considerable complexity, prohibitive for a direct numerical application, especially due to high dimensional integrals. We, therefore, will have to proceed to a quite intensive study of vertices, followed by reasonable approximations, in order to reduce drastically the numerical difficulty of the expressions.

A first purely formal simplification which we will introduce, is to consider instead of the quartet case, only the trion case. Trions are fermions and one would have to develop a whole proper philosophy to introduce an order parameter for trions. However, we will ignore this difficulty here, proceed formally as if a trion order parameter existed in the same way as a quartet order parameter, and explain for this much simpler case our strategy. This can be done without loss of generality and in the end we simply will give our results for the quartet case which can be derived in complete analogy to the trion case.

Let us, therefore, begin with the expression of the single particle mass operator containing a 'trion condensate' which analogously to Eq. (B26) is given by

$$M_{1;1'}(\omega) = \Gamma^{(3)}_{123;456}(\omega) \langle c_6 c_5 c_4 \rangle \frac{\bar{f}_2 \bar{f}_3 - f_2 f_3}{\omega + \varepsilon_{23}} (\delta_{22'} \delta_{33'} - \delta_{23'} \delta_{32'}) \times \langle c^\dagger_{4'} c^\dagger_{5'} c^\dagger_{6'} \rangle \Gamma^{(3)*}_{1'2'3';4'5'6'}(\omega) \quad \text{(D1)}$$



FIG. 11: The graphical expression of the first term on the r.h.s. of (11).



FIG. 12: Topologically same diagram as in Fig. 11.

with $\Gamma^{(3)}$ the three body vertex already given in (11) and derived in Eq. (B13). The first and rather obvious approximation we perform is to make $\Gamma^{(3)}$ 'instantaneous', that is ω-independent. A standard procedure for this is to put the vertex 'on the energy shell'. This procedure is not always defined unambiguously. In the present case, one possibility certainly is to put in $\Gamma^{(3)}$, $\omega = -\varepsilon_{23}$, i.e. the energy where the mass operator (D1) is resonant.

Let us for the moment only investigate the first term on the r.h.s. of (11) with this prescription. We obtain

$$\Gamma^{(3,1)}_{123;456}(\omega = -\varepsilon_{23}) = \frac{1}{4}\bar{v}_{13,3'6}\frac{1}{-\varepsilon_2 - \varepsilon_{3'} - \varepsilon_6}\bar{v}_{3'2,45} \qquad \text{(D2)}$$

A graphical interpretation of this term is given in Fig. 11. This graph also can be interpreted as a particular second order term of a three body scattering process (123)→(456), graphically represented in Fig. 12.
The intermediate propagator between the two vertices is given by $\frac{1}{-\varepsilon_2-\varepsilon_{3'}-\varepsilon_6}$ (the energies are given by the propagators which are cut by the vertical line in Fig. 12). We, therefore, see that the static, on-shell part $\Gamma^{(3)}(\omega = -\varepsilon_{23})$ of (D1) can be interpreted as a second order three body scattering taken at $\omega = 0$, i.e. in reality at three times the Fermi energy. This certainly is a reasonable reduction of the second order process to a static vertex $\Gamma^{(3,1)}$.

Proceeding in the same way with the second term on r.h.s. of (11), we arrive at a second order process analogous to the one of Fig. 12 as shown in Fig. 13. We see that Fig. 13 corresponds to Fig. 12 with particles 2 and 3 permuted.

Let us now consider the third term in (11). In analogy to processes of Figs. 12 and 13 this corresponds to the graph of Fig. 14.



FIG. 13: The graphical expression of the second term on the r.h.s. of (11).



FIG. 14: The graphical expression of the third term on the r.h.s. of (11).

If we want to interpret this third term also as a static second order three body process with an intermediate propagator $\frac{1}{-\varepsilon_1-\varepsilon_{2'}-\varepsilon_6}$ at $\omega = 0$, contrary to first and second terms of (11), we have to use as on shell prescription in (11) $\omega = \varepsilon_1 + \varepsilon_4 + \varepsilon_5 + \varepsilon_6$. We, thus, suppose from now on that all vertices in $\Gamma^{(3)}$ are obtained from a second order three body scattering process at $\omega = 0$.

After these preliminary considerations, let us now introduce in a phenomenological intuitive way a single particle mass operator containing a trion order parameter in a way analogous to the pairing case

$$M_{1;1'}(\omega) = \int \frac{d\omega'}{2\pi}\mathrm{Im}T_{123;1'2'3'}(\omega - \omega')G^0_{23,2'3'}(-\omega'), \qquad \text{(D3)}$$

with

$$\mathrm{Im}T_{123;1'2'3'}(\omega) = -2\pi\delta(\omega)\bar{V}_{123;456}\psi^t_{456}\psi^{t*}_{4'5'6'}\bar{V}_{4'5'6';1'2'3'}, \qquad \text{(D4)}$$

and

$$\bar{V}_{123;1'2'3'} = \frac{1}{2}(\bar{v}_{12,1'2'}\delta_{33'} + \bar{v}_{13,1'3'}\delta_{22'} + \bar{v}_{23,2'3'}\delta_{11'}), \qquad \text{(D5)}$$

where we abbreviated the the order parameter as

$$\langle c_3 c_2 c_1\rangle = \psi^t_{123}. \qquad \text{(D6)}$$

The 'three body T-matrix' (D4) in (D3) is contracted with the antisymmetrized two hole propagator

$$G^0_{23,2'3'}(-\omega) = -\frac{\bar{f}_2\bar{f}_3 - f_2 f_3}{\omega + \varepsilon_{23}}(\delta_{22'}\delta_{33'} - \delta_{23'}\delta_{32'}). \qquad \text{(D7)}$$

With the above definitions (D3)-(D7) we indeed see that the 'trion' mass operator is constructed in full analogy to the one of the pairing case in Eq. (2). It is evident

that in the quartet case we would define a four-body T-matrix as

$$\mathrm{Im}T_{1234;1'2'3'4'}(\omega) = -2\pi\delta(\omega)\Delta_{1234}\Delta^*_{1'2'3'4'} \quad \text{(D8)}$$

with

$$\Delta_{1234;5678} = \bar{V}_{1234;5678}\langle c_8 c_7 c_6 c_5\rangle \quad \text{(D9)}$$

and

$$\begin{aligned}\bar{V}_{1234;1'2'3'4'} &= \frac{1}{2}\bar{v}_{12;1'2'}\delta_{33'}\delta_{44'} + \frac{1}{2}\bar{v}_{13;1'3'}\delta_{22'}\delta_{44'} \\ &\quad +\text{permutations}\end{aligned} \quad \text{(D10)}$$

and contract it with a 3h GF.

Let us now investigate whether we can make contact of the mass operator in (D3) with the one of (D1). To this purpose, let us use in (D4) the equation for the trion order parameter which in analogy to (C3) is given by

$$\begin{aligned}\psi^t_{123} &= -\frac{1}{\varepsilon_{123}}\Big[(1-\rho_1-\rho_2)\frac{1}{2}\bar{v}_{12,1'2'}\delta_{33'} \\ &\quad +(1-\rho_1-\rho_3)\frac{1}{2}\bar{v}_{13,1'3'}\delta_{22'} \\ &\quad +(1-\rho_2-\rho_3)\frac{1}{2}\bar{v}_{23,2'3'}\delta_{11'}\Big]\psi^t_{1'2'3'}\end{aligned} \quad \text{(D11)}$$

and investigate for the moment only the expression corresponding to the second term on the r.h.s. of (D5). We obtain

$$\begin{aligned}&\frac{1}{2}\bar{v}_{13,46}\delta_{25}\psi^t_{456} \\ &= \frac{1}{2}\bar{v}_{13,46}\delta_{25}\frac{1}{-\varepsilon_{123}}\Big[(1-\rho_4-\rho_5)\frac{1}{2}\bar{v}_{45,4'5'}\delta_{66'} \\ &\quad +(1-\rho_4-\rho_6)\frac{1}{2}\bar{v}_{46,4'6'}\delta_{55'} \\ &\quad +(1-\rho_5-\rho_6)\frac{1}{2}\bar{v}_{56,5'6'}\delta_{44'}\Big]\psi^t_{4'5'6'}.\end{aligned} \quad \text{(D12)}$$

We notice some similarity with the second order vertex (D1). The difference stems from the presence of the occupation numbers ρ_i in (D12) and from the fact that there is more than one term. In principle the occupation numbers are the correlated ones. In the main part of the paper, we have seen that our theory practically only is valid for $\mu < 0$ implying that the occupation numbers remain small, see Fig. 6. We, therefore, can neglect the occupation numbers in (D12) to good approximation (for the $\mu < 0$ regime). Then, we can read off a vertex from (D12) of the following form

$$\begin{aligned}&\tilde{\Gamma}^{(3,2)}_{123;456} \\ &= \frac{1}{4}\bar{v}_{13,3'6}\frac{1}{-\varepsilon_{23'6}}\bar{v}_{3'2,45} + \frac{1}{4}\bar{v}_{13,3'6'}\delta_{25}\frac{1}{-\varepsilon_{23'6'}}\bar{v}_{3'6',46} \\ &\quad +\frac{1}{4}\bar{v}_{13,46'}\frac{1}{-\varepsilon_{246'}}\bar{v}_{26',56}\end{aligned} \quad \text{(D13)}$$

It is realized that the first term is equal to expression (D1). The second term is disconnected and, thus, is an inproper term for a vertex (see discussion below). The third term can be graphically represented as shown in Fig. 15. We see that the term of Fig. 15 is obtained from Fig. 12 in permuting indices 4 and 6. It is thus equivalent to the first term in (D13). This game can be repeated in the same way for the first and third term in (D5). The conclusion is always the same: there are twice as many terms as equivalent terms from the on shell version of (11) and in addition each time, there appears a disconnected term which should not be present in a vertex. This latter problem is, however, not a real one. The order parameter is fully correlated and no disconnected pieces can appear. The term which is disconnected in (D12) will certainly become connected in higher orders. Therefore such disconnected terms only serve to renormalize the vertices of the connected terms. In the end all bare vertices in the connected diagrams should be replaced by two body T-matrices and no disconnected terms would appear anymore. We will not further dwell on this point and simply discard the disconnected terms.



FIG. 15: The graphical expression of the third term on the r.h.s. of (D13).

The conclusion, therefore, is that from (D3) we get twice as many terms for the vertex $\Gamma^{(3)}$ as is given in Eq. (11). Otherwise the terms are equivalent under the condition that in (11) we take the above discussed on shell prescription. The correct procedure, therefore, is to divide the vertices $V_{123;1'2'3'}$ in (D4) by a factor two. In the quartet case we have to divide the vertex in (D10) by a factor of 8.

In our numerical application (quartet case), instead of calculating all the 36 terms, where several ones give equal contributions, resulting from squaring the vertex (D10) where some are more difficult to calculate than others, we take a very pragmatic point of view and keep only the first term on the r.h.s. of (D10) simulating all the others by a factor λ' as in Eq. (18). In doing so, we suppose that all terms have more or less the same analytic structure. This is certainly the case, since all the terms are dominated by the behavior of the 3h level density whose typical structure is displayed in Sec. V. It also is fortunate that the final result does not depend on λ', as we now will demonstrate on the example of pairing.

Appendix E: Independence of the parameter λ' in Eq. (18)

All arguments and derivations for the three-body vertices can directly be generalized to the four body case.

In view of the fact that our favorite expression (8) is too complex for a direct numerical realization, we take a very pragmatic point of view for a pilot application. We have seen that some terms appear in all three forms we discussed for the vertices. We shall take one of those terms for numerical calculations with a parameter λ' in front of the vertex which should mock up the influence of factors and also of topologically different graphs. However, we want to stress again that we do not think that topologically different vertices will finally give rise to different analytic structure of the mass operator. Essentially these additional terms will again only renormalize the vertices. The form which we then retain is just the first one on the r.h.s. of (D10), since it is the one where the single particle motion directly couples via an interaction to the quartet amplitude, that is

$$\tilde{\Delta}_{1234} = \lambda' \frac{1}{2} \bar{v}_{12,1'2'} \delta_{33'} \delta_{44'} \langle c_{1'} c_{2'} c_{3'} c_{4'} \rangle \tag{E1}$$

With the separable form of the two body interaction [21], we then obtain expression (21). As mentioned already, we will approximately absorb factors and all the other terms in renormalizing the vertex by a constant factor λ'. This shall be a quite valid procedure, since the analytic structure of all the other terms is very similar. Fortunately our results will not depend on the strength of the renormalization factor λ' of the vertex. This statement may be surprising but can be explained at hand of the standard BCS example as follows.

Schematically, we write expression for single particle GF in BCS approximation of Eqs. (1) and (2) as follows (with self evident notation)

$$G^{\rm BCS} = \frac{1}{\omega - \varepsilon - \dfrac{v\langle cc\rangle\langle c^\dagger c^\dagger\rangle v}{\omega + \varepsilon}} \tag{E2}$$

Now let us change in above equation vertex v to $v' = \lambda' v$. Let us apply to all quantities containing this new vertex a 'prime'. Then BCS single particle GF is given by

$$G'^{\rm BCS} = \frac{1-\rho'}{\omega - E'} + \frac{\rho'}{\omega + E'} \tag{E3}$$

with $E'^2 = \varepsilon^2 + D'^2$, $D' = v'\langle cc\rangle$, and $\rho' = (1 - \varepsilon/E')/2$. The equation for order parameter then reads

$$\langle cc\rangle = -\frac{1-2\rho'}{2\varepsilon} v \langle cc\rangle \tag{E4}$$

where we should pay attention that the v in that equation is still the original one, as in our four-body order parameter equation of (C2) and (C5).

From above we obtain the gap equation for Δ'

$$\Delta' = -\sum v \frac{\Delta'}{2E'} \tag{E5}$$

Multiplying this equation with λ', we see that $E' = E$ and, therefore, nothing is changed in multiplying in BCS single particle mass operator the vertex by an arbitrary factor. Translated to our quartet problem, this implies that the question of multiplicity of diagrams discussed above is of no consequence for our numerical results. Of course, the addition of topologically different contributions to the vertex $\Gamma^{(4)}$ can change things slightly but again one can imagine that to good approximation it only renormalizes the vertex corresponding to the process which we treat explicitly. Therefore, our approximate treatment of the single particle mass operator with quartet condensate should be quite safe. With our separable ansatz for the two body interaction, we then arrive to expression (B20) of the mass operator.

Appendix F: Preparation of mass operator for numerical application

In this section, we describe the final expression for numerical calculation of the mass operator of Eq. (21).

We again give the approximated mass operator

$$\begin{aligned}
& M^{\rm quartet}(k_1,\omega) \\
= & \int \frac{d^3k_2}{(2\pi)^3}\frac{d^3k_3}{(2\pi)^3}\frac{d^3k_4}{(2\pi)^3}\frac{d^3k_{1'}}{(2\pi)^3}\frac{d^3k_{2'}}{(2\pi)^3}\frac{d^3k_{1''}}{(2\pi)^3}\frac{d^3k_{2''}}{(2\pi)^3} \\
\times & \; e^{-2(\vec{k}_1-\vec{k}_2)^2/(4b^2)} e^{-(\vec{k}_{1'}-\vec{k}_{2'})^2/(4b^2)} e^{-(\vec{k}_{1''}-\vec{k}_{2'''})^2/(4b^2)} (2\pi)^3 \delta(\vec{k}_1+\vec{k}_2-\vec{k}_{1'}-\vec{k}_{2'}) \\
\times & \; \varphi(|\vec{k}_{1'}|)\varphi(|\vec{k}_{2'}|)\varphi(|\vec{k}_3|)\varphi(|\vec{k}_4|)(2\pi)^3\delta(\vec{k}_{1'}+\vec{k}_{2'}+\vec{k}_3+\vec{k}_4) \\
\times & \; \varphi(|\vec{k}_{1''}|)\varphi(|\vec{k}_{2''}|)\varphi(|\vec{k}_3|)\varphi(|\vec{k}_4|)(2\pi)^3\delta(\vec{k}_{1''}+\vec{k}_{2''}+\vec{k}_3+\vec{k}_4) \\
\times & \; \frac{\bar{f}(k_2)\bar{f}(k_3)\bar{f}(k_4) + f(k_2)f(k_3)f(k_4)}{\omega + \varepsilon_{\vec{k}_2} + \varepsilon_{\vec{k}_3} + \varepsilon_{\vec{k}_4}} \\
= & \int \frac{d^3k_2}{(2\pi)^3}\frac{d^3K}{(2\pi)^3}\frac{d^3k}{(2\pi)^3}\frac{d^3K'}{(2\pi)^3}\frac{d^3k'}{(2\pi)^3}\frac{d^3K''}{(2\pi)^3}\frac{d^3k''}{(2\pi)^3}
\end{aligned}$$

$$
\begin{aligned}
&\times\ e^{-2(\vec{k}_1-\vec{k}_2)^2/(4b^2)}e^{-k'^2/b^2}e^{-k''^2/b^2}(2\pi)^3\delta(\vec{k}_1+\vec{k}_2-\vec{K}')\\
&\times\ \varphi(|\frac{\vec{K}'}{2}+\vec{k}'|)\varphi(|\frac{\vec{K}'}{2}-\vec{k}'|)\varphi(|\frac{\vec{K}}{2}+\vec{k}|)\varphi(|\frac{\vec{K}}{2}-\vec{k}|)(2\pi)^3\delta(\vec{K}'+\vec{K})\\
&\times\ \varphi(|\frac{\vec{K}''}{2}+\vec{k}''|)\varphi(|\frac{\vec{K}''}{2}-\vec{k}''|)\varphi(|\frac{\vec{K}}{2}+\vec{k}|)\varphi(|\frac{\vec{K}}{2}-\vec{k}|)(2\pi)^3\delta(\vec{K}''+\vec{K})\\
&\times\ \frac{\bar{f}(k_2)\bar{f}(|\frac{\vec{K}}{2}+\vec{k}|)\bar{f}(|\frac{\vec{K}}{2}-\vec{k}|)+f(k_2)f(|\frac{\vec{K}}{2}+\vec{k}|)f(|\frac{\vec{K}}{2}-\vec{k}|)}{\omega+\varepsilon_{\vec{k}_2}+\varepsilon_{\frac{\vec{K}}{2}+\vec{k}}+\varepsilon_{\frac{\vec{K}}{2}-\vec{k}}}\\
&=\ \int\frac{d^3K}{(2\pi)^3}\frac{d^3k}{(2\pi)^3}e^{-2(\vec{k}_1+\frac{\vec{K}}{2})^2/b^2}\Big[\varphi(|\frac{\vec{K}}{2}+\vec{k}|)\varphi(|\frac{\vec{K}}{2}-\vec{k}|)\Big]^2\\
&\times\ \frac{\bar{f}(|\vec{k}_1+\vec{K}|)\bar{f}(|\frac{\vec{K}}{2}+\vec{k}|)\bar{f}(|\frac{\vec{K}}{2}-\vec{k}|)+f(|\vec{k}_1+\vec{K}|)f(|\frac{\vec{K}}{2}+\vec{k}|)f(|\frac{\vec{K}}{2}-\vec{k}|)}{\omega+\frac{k_1^2}{2m}+\frac{\vec{k}_1\cdot\vec{K}}{m}+\frac{3K^2}{4m}+\frac{k^2}{m}-3\mu}\\
&\times\ \Big[\int\frac{d^3k'}{(2\pi)^3}e^{-k'^2/b^2}\varphi(|-\frac{\vec{K}}{2}+\vec{k}'|)\varphi(|-\frac{\vec{K}}{2}-\vec{k}'|)\Big]^2. \qquad \text{(F1)}
\end{aligned}
$$

The imaginary part of the above mass operator $-\mathrm{Im}M^{\mathrm{quartet}}(k_1,\omega+i\eta)$ is

$$
\begin{aligned}
&-\mathrm{Im}M^{\mathrm{quartet}}(k_1,\omega+i\eta)\\
&=\ \pi\int\frac{d^3K}{(2\pi)^3}\frac{d^3k}{(2\pi)^3}e^{-2(\vec{k}_1+\frac{\vec{K}}{2})^2/b^2}\\
&\times\ \Big[\varphi(|\frac{\vec{K}}{2}+\vec{k}|)\varphi(|\frac{\vec{K}}{2}-\vec{k}|)\Big]^2\\
&\times\ \Big[\bar{f}(|\vec{k}_1+\vec{K}|)\bar{f}(|\frac{\vec{K}}{2}+\vec{k}|)\bar{f}(|\frac{\vec{K}}{2}-\vec{k}|)\\
&\quad+f(|\vec{k}_1+\vec{K}|)f(|\frac{\vec{K}}{2}+\vec{k}|)f(|\frac{\vec{K}}{2}-\vec{k}|)\Big]\\
&\times\ \delta(\omega+\frac{k_1^2}{2m}+\frac{\vec{k}_1\cdot\vec{K}}{m}+\frac{3K^2}{4m}+\frac{k^2}{m}-3\mu)\\
&\times\ \Big[\int\frac{d^3k'}{(2\pi)^3}e^{-k'^2/b^2}\varphi(|-\frac{\vec{K}}{2}+\vec{k}'|)\varphi(|-\frac{\vec{K}}{2}-\vec{k}'|)\Big]^2
\end{aligned}
\qquad \text{(F2)}
$$

1. The $\vec{k}_1=0$ case

The imaginary part of the mass operator for $\vec{k}_1=0$ is given by

$$
\begin{aligned}
&-\mathrm{Im}M^{\mathrm{quartet}}(k_1=0,\omega+i\eta)\\
&=\ \pi\int\frac{d^3K}{(2\pi)^3}\frac{d^3k}{(2\pi)^3}e^{-K^2/(4b^2)}\\
&\times\ \Big[\varphi(|\frac{\vec{K}}{2}+\vec{k}|)\varphi(|\frac{\vec{K}}{2}-\vec{k}|)\Big]^2\\
&\times\ \Big[\bar{f}(K)\bar{f}(|\frac{\vec{K}}{2}+\vec{k}|)\bar{f}(|\frac{\vec{K}}{2}-\vec{k}|)\\
&\quad+f(K)f(|\frac{\vec{K}}{2}+\vec{k}|)f(|\frac{\vec{K}}{2}-\vec{k}|)\Big]\\
&\times\ \delta(\omega+\frac{3K^2}{4m}+\frac{k^2}{m}-3\mu)\\
&\times\ \Big[\int\frac{d^3k'}{(2\pi)^3}e^{-k'^2/b^2}\varphi(|-\frac{\vec{K}}{2}+\vec{k}'|)\varphi(|-\frac{\vec{K}}{2}-\vec{k}'|)\Big]^2\\
&=\ \pi\frac{2}{(2\pi)^4}\int_0^\infty dKK^2\int_0^\infty dkk^2\int_{-1}^1 dte^{-K^2/(4b^2)}\\
&\times\ \Big[\varphi(|\frac{\vec{K}}{2}+\vec{k}|)\varphi(|\frac{\vec{K}}{2}-\vec{k}|)\Big]^2\\
&\times\ \Big[\bar{f}(K)\bar{f}(|\frac{\vec{K}}{2}+\vec{k}|)\bar{f}(|\frac{\vec{K}}{2}-\vec{k}|)\\
&\quad+f(K)f(|\frac{\vec{K}}{2}+\vec{k}|)f(|\frac{\vec{K}}{2}-\vec{k}|)\Big]\\
&\times\ \frac{m}{2k}\delta(k-\sqrt{3m\mu-m\omega-3K^2/4})\\
&\times\ \Big[\frac{1}{(2\pi)^2}\int_0^\infty dk'k'^2\int_{-1}^1 dt'e^{-k'^2/b^2}\\
&\times\ \varphi(|-\frac{\vec{K}}{2}+\vec{k}'|)\varphi(|-\frac{\vec{K}}{2}-\vec{k}'|)\Big]^2\\
&=\ \frac{2\pi m}{2(2\pi)^4}\int_0^{P_{max0}} dKK^2p\int_{-1}^1 dte^{-K^2/(4b^2)}\\
&\times\ \Big[\varphi(|\frac{\vec{K}}{2}+\vec{p}|)\varphi(|\frac{\vec{K}}{2}-\vec{p}|)\Big]^2\\
&\times\ \Big[\bar{f}(K)\bar{f}(|\frac{\vec{K}}{2}+\vec{p}|)\bar{f}(|\frac{\vec{K}}{2}-\vec{p}|)\\
&\quad+f(K)f(|\frac{\vec{K}}{2}+\vec{p}|)f(|\frac{\vec{K}}{2}-\vec{p}|)\Big]\\
&\times\ \Big[\frac{1}{(2\pi)^2}\int_0^\infty dk'k'^2\int_{-1}^1 dt'e^{-k'^2/b^2}\\
&\times\ \varphi(|-\frac{\vec{K}}{2}+\vec{k}'|)\varphi(|-\frac{\vec{K}}{2}-\vec{k}'|)\Big]^2,
\end{aligned}
\qquad \text{(F3)}
$$

Here, in the last equation,

$$p = \sqrt{3m\mu - m\omega - \frac{3K^2}{4}}, \tag{F4}$$

$$P_{max0} = \sqrt{\frac{4}{3}(3m\mu - m\omega)}, \tag{F5}$$

$$t = (\vec{p}\cdot\vec{K})/(pK), \tag{F6}$$

$$t' = (\vec{k}'\cdot\vec{K})/(k'K). \tag{F7}$$

2. The $\vec{k}_1 \neq 0$ case

On the other hand, $-\mathrm{Im}M^{\mathrm{quartet}}$ for $\vec{k}_1 \neq 0$ is represented by

$$\begin{aligned}
&-\mathrm{Im}M^{\mathrm{quartet}}(k_1, \omega + i\eta) \\
&= \frac{\pi}{(2\pi)^4}\int_0^\infty dKK^2\int_{-1}^1 dt_1\int_0^\infty dkk^2\int_{-1}^1 dt \\
&\times\ e^{-2(k_1^2+K^2/4+k_1Kt_1)/b^2}\Big[\varphi(|\frac{\vec{K}}{2}+\vec{k}|)\varphi(|\frac{\vec{K}}{2}-\vec{k}|)\Big]^2 \\
&\times\ \Big[\bar{f}(|\vec{k}_1+\vec{K}|)\bar{f}(|\frac{\vec{K}}{2}+\vec{k}|)\bar{f}(|\frac{\vec{K}}{2}-\vec{k}|) \\
&\quad + f(|\vec{k}_1+\vec{K}|)f(|\frac{\vec{K}}{2}+\vec{k}|)f(|\frac{\vec{K}}{2}-\vec{k}|)\Big] \\
&\times\ \delta(\omega + \frac{k_1^2}{2m} + \frac{k_1Kt_1}{m} + \frac{3K^2}{4m} + \frac{k^2}{m} - 3\mu) \\
&\times\ \Big[\frac{1}{(2\pi)^2}\int dk'k'^2\int_{-1}^1 dt' e^{-k'^2/b^2} \\
&\times\ \varphi(|-\frac{\vec{K}}{2}+\vec{k}'|)\varphi(|-\frac{\vec{K}}{2}-\vec{k}'|)\Big]^2 \\
&= \frac{\pi}{(2\pi)^4}\int_0^\infty dKK^2\int_{-1}^1 dt_1\int_0^\infty dkk^2\int_{-1}^1 dt \\
&\times\ e^{-2(k_1^2+K^2/4+k_1Kt_1)/b^2}\Big[\varphi(|\frac{\vec{K}}{2}+\vec{k}|)\varphi(|\frac{\vec{K}}{2}-\vec{k}|)\Big]^2 \\
&\times\ \Big[\bar{f}(|\vec{k}_1+\vec{K}|)\bar{f}(|\frac{\vec{K}}{2}+\vec{k}|)\bar{f}(|\frac{\vec{K}}{2}-\vec{k}|) \\
&\quad + f(|\vec{k}_1+\vec{K}|)f(|\frac{\vec{K}}{2}+\vec{k}|)f(|\frac{\vec{K}}{2}-\vec{k}|)\Big] \\
&\times\ \frac{m}{k_1K}\delta(t_1 - \frac{3m\mu - m\omega - \frac{k_1^2}{2} - \frac{3K^2}{4} - k^2}{k_1K}) \\
&\times\ \Big[\frac{1}{(2\pi)^2}\int dk'k'^2\int_{-1}^1 dt' e^{-k'^2/b^2} \\
&\times\ \varphi(|-\frac{\vec{K}}{2}+\vec{k}'|)\varphi(|-\frac{\vec{K}}{2}-\vec{k}'|)\Big]^2.
\end{aligned} \tag{F8}$$

Here, the following condition has to be satisfied:

$$-1 < \frac{3m\mu - m\omega - \frac{k_1^2}{2} - \frac{3K^2}{4} - k^2}{k_1K} < 1. \tag{F9}$$

Therefore

$$k^2 > 3m\mu - m\omega - \frac{k_1^2}{2} - \frac{3K^2}{4} - k_1K > 0 \tag{F10}$$

and

$$0 < k^2 < 3m\mu - m\omega - \frac{k_1^2}{2} - \frac{3K^2}{4} + k_1K. \tag{F11}$$

From the above equation, we obtain the limits of integration with respect to k:

$$\begin{aligned}
p_{min}(K) &= \sqrt{\max(3m\mu - m\omega - \frac{k_1^2}{2} - \frac{3K^2}{4} - k_1K, 0)} \\
&< k < \sqrt{3m\mu - m\omega - \frac{k_1^2}{2} - \frac{3K^2}{4} + k_1K} = p_{max}(K)
\end{aligned} \tag{F12}$$

Besides, from Eq. (F11),

$$3m\mu - m\omega - \frac{k_1^2}{2} - \frac{3K^2}{4} + k_1K > 0 \tag{F13}$$

has to be satisfied. Therefore

$$\begin{aligned}
P_{min} &= \max\left[\frac{2}{3}\Big(k_1 - \sqrt{9m\mu - 3m\omega - \frac{k_1^2}{2}}\Big), 0\right] \\
&< K < \frac{2}{3}\left(k_1 + \sqrt{9m\mu - 3m\omega - \frac{k_1^2}{2}}\right) = P_{max}
\end{aligned} \tag{F14}$$

3. Summary of integrals in $\mathrm{Im}M^{\mathrm{quartet}}$

With the Appendices F 1 and F 2, we can express $-\mathrm{Im}M^{\mathrm{quartet}}(k_1, \omega + i\eta)$ by:

If $9m\mu - 3m\omega - \frac{k_1^2}{2} < 0$,

$$-\mathrm{Im}M^{\mathrm{quartet}}(k_1, \omega + i\eta) = 0. \tag{F15}$$

Else:

If $k_1 = 0$,

$$\begin{aligned}
&-\mathrm{Im}M^{\mathrm{quartet}}(k_1, \omega + i\eta) \\
&= \frac{2\pi m}{2(2\pi)^4}\int_0^{P_{max0}} dKK^2pe^{-K^2/(4b^2)} \\
&\times\ \int_{-1}^1 dt\left[\varphi(|\frac{\vec{K}}{2}+\vec{p}|)\varphi(|\frac{\vec{K}}{2}-\vec{p}|)\right]^2 \\
&\times\ \Big[\bar{f}(K)\bar{f}(|\frac{\vec{K}}{2}+\vec{p}|)\bar{f}(|\frac{\vec{K}}{2}-\vec{p}|) \\
&\quad + f(K)f(|\frac{\vec{K}}{2}+\vec{p}|)f(|\frac{\vec{K}}{2}-\vec{p}|)\Big]
\end{aligned}$$

$$\times \Big[\frac{1}{(2\pi)^2}\int_0^\infty dk' k'^2 e^{-k'^2/b^2}\int_{-1}^1 dt' \times \varphi(|-\frac{\vec{K}}{2}+\vec{k}'|)\varphi(|-\frac{\vec{K}}{2}-\vec{k}'|)\Big]^2,$$

$$p=\sqrt{3m\mu-m\omega-\frac{3K^2}{4}},$$

$$P_{max0}=\sqrt{\frac{4}{3}(3m\mu-m\omega)},$$

$$t=(\vec{k}\cdot\vec{K})/(kK), \qquad t'=(\vec{k}'\cdot\vec{K})/(k'K) \quad \text{(F16)}$$

while if $k_1 \neq 0$,

$$\begin{aligned}
&-\mathrm{Im}M^{\mathrm{quartet}}(k_1,\omega+i\eta)\\
&= \frac{\pi m}{(2\pi)^4 k_1}\int_{P_{min}}^{P_{max}} dKK\int_{p_{min}}^{p_{max}} dkk^2\\
&\times e^{-2(3m\mu-m\omega+k_1^2/2-K^2/2-k^2)/b^2}\\
&\times \int_{-1}^1 dt\Big[\varphi(|\frac{\vec{K}}{2}+\vec{k}|)\varphi(|\frac{\vec{K}}{2}-\vec{k}|)\Big]^2\\
&\times \Big[\bar{f}(\sqrt{6m\mu-2m\omega-\frac{K^2}{2}-2k^2})\bar{f}(|\frac{\vec{K}}{2}+\vec{k}|)\bar{f}(|\frac{\vec{K}}{2}-\vec{k}|)\\
&\quad +f(\sqrt{6m\mu-2m\omega-\frac{K^2}{2}-2k^2})f(|\frac{\vec{K}}{2}+\vec{k}|)f(|\frac{\vec{K}}{2}-\vec{k}|)\Big]\\
&\times \Big[\frac{1}{(2\pi)^2}\int_0^\infty dk' k'^2 e^{-k'^2/b^2}\int_{-1}^1 dt' \times \varphi(|-\frac{\vec{K}}{2}+\vec{k}'|)\varphi(|-\frac{\vec{K}}{2}-\vec{k}'|)\Big]^2,
\end{aligned}$$

$$p_{min}(K)=\sqrt{\max[3m\mu-m\omega-\frac{k_1^2}{2}-\frac{3K^2}{4}-k_1K,0]},$$

$$p_{max}(K)=\sqrt{3m\mu-m\omega-\frac{k_1^2}{2}-\frac{3K^2}{4}+k_1K},$$

$$P_{min}=\max\left[\frac{2}{3}\left(k_1-\sqrt{9m\mu-3m\omega-\frac{k_1^2}{2}}\right),0\right],$$

$$P_{max}=\frac{2}{3}\left(k_1+\sqrt{9m\mu-3m\omega-\frac{k_1^2}{2}}\right),$$

$$t=(\vec{k}\cdot\vec{K})/(kK), \qquad t'=(\vec{k}'\cdot\vec{K})/(k'K). \quad \text{(F17)}$$

$\mathrm{Re}M^{\mathrm{quartet}}$ is calculated by Eq. (22).

[1] A. Tohsaki, H. Horiuchi, P. Schuck, and G. Röpke, Phys. Rev. Lett. **87**, 192501 (2001); Y. Funaki, T. Yamada, H. Horiuchi, G. Röpke, P. Schuck, and A. Tohsaki, Phys. Rev. Lett. **101**, 082502 (2008).

[2] S. Typel1, G. Röpke, T. Klähn, D. Blaschke, and H. H. Wolter, Phys. Rev. C **81**, 015803 (2010).

[3] A. N. Wenz, T. Lompe, T. B. Ottenstein, F. Serwane, G. Zörn, and S. Jochim, Phys. Rev. A **80**, 040702(R) (2009).

[4] J. R. Williams, E. L. Hazlett, J. H. Huckans, R. W. Stites, Y. Zhang, and K. M. O'Hara, Phys. Rev. Lett. **103**, 130404 (2009).

[5] Ákos Rapp, Gergely Zaránd, Carsten Honerkamp, and Walter Hofstetter, Phys. Rev. Lett. **98**, 160405 (2007).

[6] X. W. Guan, M. T. Batchelor, C. Lee, and H.-Q. Zhou Phys. Rev. Lett. **100**, 200401 (2008).

[7] A. Kantian, M. Dalmonte, S. Diehl, W. Hofstetter, P. Zoller, and A. J. Daley, Phys. Rev. Lett. **103**, 240401 (2009).

[8] S. Floerchinger, R. Schmidt, and C. Wetterich, Phys. Rev. A **79**, 053633 (2009).

[9] Beatriz Errea, Jorge Dukelsky, and Gerardo Ortiz, Phys. Rev. A **79**, 051603 (2009).

[10] Theja N. De Silva, Phys. Rev. A **80**, 013620 (2009).

[11] Rafael A. Molina, Jorge Dukelsky, and Peter Schmitteckert, Phys. Rev. A **80**, 013616 (2009).

[12] P. Azaria, S. Capponi, and P. Lecheminant, Phys. Rev. A **80**, 041604 (2009).

[13] Shin-ya Miyatake, Kensuke Inaba, and Sei-ichiro Suga, Phys. Rev. A **81**, 021603 (2010).

[14] H. Kamei and K. Miyake, J. Phys. Soc. Jpn. **74**, 1911 (2005).

[15] A. S. Stepanenko and J. M. Gunn, arXiv:cond-mat/9901317.

[16] P. Schlottmann, J. Phys. Condens. Matter **6**, 1359 (1994).

[17] C. Wu, Phys. Rev. Lett. **95**, 266404 (2005).

[18] G. Roux, S. Capponi, P. Lecheminant, and P. Azaria, Eur. Phys. J. B **68**, 293 (2009).

[19] M. Hasuo and N. Nagasawa, in *Bose-Einstein Condensation* edited by A. Greffin, D. W. Snoke and S. Stringari (Cambridge University Press, Cambridge, 1995), p. 487.

[20] S. A. Moskalenko and D. W. Snoke, *Bose-Einstein Condensation of Excitons and Biexcitons* (Cambridge University Press, Cambridge, 2000).

[21] T. Sogo, R. Lazauskas, G. Röpke, and P. Schuck, Phys. Rev. C **79**, 051301 (2009).

[22] G. Röpke, A. Schnell, P. Schuck, and P. Nozières, Phys. Rev. Lett. **80**, 3177 (1998).

[23] D. J. Thouless, Ann. Phys. **10**, 553 (1960).

[24] A. A. Abrikosov, L. P. Gorkov, and I. E. Dzyaloshinski, *Mothods of quantum field theory in statistical physics*, (Dover, New York, 1975).

[25] A. L. Fetter and J. D. Walecka, *Quantum Theory of Many-Particle Systems*, (Dover, New York, 2003).

[26] P. Ring and P. Schuck, *The Nuclear Many-Body Problem*, (Springer-Verlag, New York, 1980)

[27] J.G. Hirsch, A. Mariano, J. Dukelsky, and P. Schuck, Ann. Phys. **296**, 187 (2002).

[28] A. H. Blin, R. W. Hasse, B. Hiller, P. Schuck, and C. Yannouleas, Nucl. Phys. **A456**, 109 (1986).

[29] S. Adachi and P. Schuck, Nucl. Phys. **A496**, 485 (1989).

[30] J. Dukelsky, G. Röpke, and P. Schuck, Nucl. Phys. **A628**, 17 (1998).

[31] G. D. Mahan, *Many-Particle Physics*, (Plenum, New York, 1990).

[32] P. Nozières and S. Schmitt-Rink, J. Low Temp. Phys. **59**,

195 (1985).
[33] G. F. Bertsch, P. F. Bortignon, and R. A. Broglia, Rev. Mod. Phys. **55**, 287 (1983).
[34] K. Ando, A. Ikeda and, G. Holzwarth, Z. Phys. **A310** 223 (1983).
[35] G. Röpke, L. Münchow, and H. Schulz, Nucl. Phys. **A379**, 536 (1982); G. Röpke, M. Schmidt, L. Münchow, and H. Schulz, Nucl. Phys. **A399**, 587 (1983).